



Available online on 15.1.2018 at <http://ujpr.org>

Universal Journal of Pharmaceutical Research

An International Peer Reviewed Journal

Open access to Pharmaceutical research

© 2017, publisher and licensee UJPR, This is an open access article which permits unrestricted non commercial use, provided the original work is properly cited

Volume 2, Issue 6, 2017

RESEARCH ARTICLE

INVESTIGATION OF TOXIC METALS POLLUTION IN WATER, SEDIMENT AND FISH AT ADEN COAST, GULF OF ADEN, YEMEN

Mohammed Kassem Othman Al-qadasy¹, Abdulla Saleh Babaqi¹, Mukhtar Mohammed Al-Abyadh², Ali Gamal Ahmed Al-kaf³

¹Department of Chemistry, Faculty of Science, Sana'a University, Sana'a-Yemen. ²Department of Chemistry, Faculty of Pharmacy, University of Science and Technology, Aden-Yemen. ³Sana'a University-Faculty of pharmacy, Yemen

ABSTRACT

This study investigates concentrations of toxic metals, Lead and Cadmium in water, sediments and fish organs (muscle, liver and gill). Collected from Aden coast, gulf of Aden in Yemen. Water samples and sediment were taken from nine locations. Four fish species were collected (*Lethrinus mahsena*; *Epinephelus areolatus*; *Thunnus tonggol* and *Sphyrna jello*) were collected from the local commercial fishermen of Aden city during winter 2011, Summer 2012 and winter 2013. Lead concentration in sea water is 0.045-0.055 mg/l and Lead concentration in Sediment 33.512-35.726 µg/g dry wt., Cadmium concentration in sea water 0.006-0.010 mg/l, Cadmium concentration in sediment 1.944-2.004 µg/g dry wt., lead concentration is the highest in most fish gill samples 0.047-0.727 µg/g dry wt., where as in muscles is the lowest 0.020-0.116 µg/g dry wt., and in liver was 0.038- 0.267 µg/g dry wt. Cadmium concentration is the highest in most fish gill samples 0.033-0.609 µg/g dry wt., where as in muscles is the lowest 0.018- 0.073 µg/g dry wt., and in liver was 0.028-0.209 µg/g dry wt. By comparing the results obtained with other data obtained from the local and international studies, in addition to, comparing the results standard levels of these metals contaminated and adopted internationally and domestically and the pollution levels in Yemen is currently within the lower limits of pollution.

Keywords: Aden, Cadmium, fish, Lead, sediments, water, Yemen.

Article Info: Received 22 November 2017; Revised 8 January; Accepted 11 January, Available online 15 January 2018



Cite this article-

Mohammed Kassem Othman Al-qadasy, Abdulla Saleh Babaqi, Mukhtar Mohammed Al-Abyadh, Ali Gamal Ahmed Al-kaf. Investigation of toxic metals pollution in water, sediment and fish at Aden coast, gulf of Aden, Yemen. Universal Journal of Pharmaceutical Research. 2017; 2(6): 42-49.

DOI: <http://doi.org/10.22270/ujpr.v2i6.R8>

Address for Correspondence:

Dr. Mohammed Qasem Al-salehi, P.O. Box 11205 Sana'a, Yemen, Tel: +967-777408835, E-mail:

alqadasy64@yahoo.com

INTRODUCTION

The pollution of aquatic systems has become a major concern worldwide¹. There are a variety of sources that will pollute aquatic systems with heavy metals. These include animal matter, wet and dry fallouts of atmospheric particulate matter and human activities. The concentration, bioavailability and toxicity of heavy metals in aquatic systems can be affected by various factors, including pH and temperature². Poor quality of surface water is caused in two ways. The pollution of surface water can either be due to point source (PS) or nonpoint source pollution (NPS). Point source pollution is mainly municipal sewage discharge and industrial wastewater loads. Municipal sewage discharge is from urban or highly residential areas, while industrial wastewater is from a variety of manufacturers³. When rainfall or irrigation water runs over land it will carry and deposit pollutants into rivers,

lakes and coastal waters. This is seen as nonpoint source pollution³. Heavy metals will be distributed between the aqueous phase and bed sediments in aquatic systems⁴. Only a small percentage of the free metal ions stay dissolved in water. The majority of the ions get deposited in the sediment due to adsorption, hydrolysis and co-precipitation of the free ions⁴.

As an important component of water environment, sediment is not only the place where pollutants accumulate from the water body, but also it is a secondary pollution source which has a potential impact on water quality⁵. Sediment represents one of ultimate sinks for heavy metals discharged into the aquatic environment. Therefore, sediment quality is a good indicator of pollution in the water column, where it tends to concentrate the heavy metals⁶. Heavy metals are distributed in sediments in four fractions, as

exchangeable bound, iron– manganese oxide, organic matter and residual species⁷.

In order to protect the aquatic life community comprehensive methods for identifying and assessing the severity of sediment contamination. Due to the ecological importance and the persistence of pollutants in the aquatic ecosystem, sediments are more appropriate to be monitored in environmental evaluations and understand their potential toxic impacts⁸. Sediment pollution, especially from heavy metals, has an important impact on the water environment and a direct potential threat on human and aquatic⁵.

Fishes represent the peak of consumers in the water system. Fishes have ability to collect these metals in concentrations higher than water and sediments because of feed on organic materials in aquatic environments⁹. Fishes have been found to be good indicators of the heavy metal contamination levels in the aquatic systems because they occupy different trophic levels¹⁰. According to there are two main routes of heavy metals exposure: The primary route of intake of these chemicals in fish species is via gill or transport of dissolved contaminants in water across biological membranes and ionic exchange. The secondary route is through the intestine by food or sediment particles with subsequent transport across the gut. The food may also be important source for heavy metal accumulation in fish¹¹. In aquatic ecosystem, metals are transferred to the fish through food chain that could ultimately affect the health of people consuming this fish. Accumulation of these metals in the bodies of fish affected by different factors such as pH, water hardness and level of pollution in the surrounding water added to the age and physiological situation of fish¹². Industrial and domestic waste containing heavy metals and hydrocarbon accumulate in aquatic food chains as possible to cause acute and chronic damages in fish communities and lead to reducibility to growth and reproduce¹³.

Lead, in water, accumulates in the body of fish and other marine organisms and it is eventually ingested by humans who consume these fish and seafood products¹⁴. The presence of Pb in the human body causes damage to the nervous system through several mechanisms. Neuropsychological research over the years has revealed that Pb exposure can result in declines in intelligence, memory, processing speed, comprehension and reading, visuospatial, motor and executive skills. Among the cognitive deficits induced by Pb toxicity, visuospatial deficits appear to be major. Anxiety, depression and phobia can also occur, while outcome, intervention, and rehabilitation results are largely dependent on the level of toxic exposure. There is also a growing evidence of antisocial behaviour linked to early Pb exposure¹⁵.

Cadmium in water can be accumulated in the body of marine organisms and can eventually enter the body of humans who consume these seafood products. The concentrations of Cd call for caution as cumulative effects might constitute health hazards to aquatic life including man who feeds on fish¹⁶. In the late 1960s environmental cadmium contamination was established

as the cause of an epidemic of bone disease (itai-itai disease) in Japan. Since that time, increasing scientific interest has been devoted to cadmium as an environmental contaminant. Awareness is now been disseminated in some countries concerning the small margin of safety between existing intake levels and levels that may cause adverse health effect to the population¹⁷.

Determining the levels of such heavy metals and comparing the levels with guidelines will establish the potential health risk from the consumption of such fish species. Therefore, it is important to determine the concentrations of non-essential metals in fish in order to evaluate the possible risks of fish consumption. This can serve as an indicator for the extent of pollution in Yemen coastal waters.

MATERIALS AND METHODS

Water samples and Sediment samples were collected from Winter 2011, Summer 2012 and Winter 2013, from 9 sampling sites (Fig. 1). during which a total of 27 Sample of Surface Seawater, 27 Sample of Sediments, 36 Sample of Muscles Fish, 36 Sample of Liver Fish and 36 Sample of Gills Fish were collected and analyzed.



Figure 1: Sampling locations (1– 9) along the Coast of Aden, Yemen.

Sea water samples were collected using cleaned plastic water sampler. Each sample was taken in 1 liter polyethylene bottles. All water samples were immediately brought to the laboratory where filtered through whatman No.41 (0.45 μm pore size) filter paper. The samples were acidified with 2ml nitric acid to prevent precipitation of metals, reduce adsorption of the analyses onto the walls of containers and to avoid microbial activity, and then stored at 4°C until the chemical analyses. Surface Seawater Digestion for Pb and Cd Analysis by GFAAS, Five milliliter of concentrated HCl was added to 250 ml of each surface seawater sample placed in 600 ml beaker and evaporated to 25 ml volume. The concentrate was transferred to a 50 ml volumetric flask and diluted to mark with deionized water. Prior analysis, the solutions

were filtered through Whatman No.41 (0.45 µm pore size) filter paper.

Analyzed for Lead (Pb) and Cadmium (Cd) using Buck Model 210 VGP, USA Made -Graphite furnace Atomic Absorption Spectrophotometer (GF AAS) in Seawater samples, before proceeding Method 200.13¹⁸

Sediments samples were collected by a polyethylene corer in accord with standard methods. The procedure for the extraction heavy metals was based on Standard Method 3051A (Microwave – assisted acid digestion of sediments¹⁹. About 0.25 g of dry sediment sample was accurately weighed and digested with 6 ml of concentrated nitric acid (HNO₃ 65%) , 1 ml of Perchloric acid (HClO₄ 65%) and 1 ml of hydrogen peroxide (H₂O₂ 30%) . Milestone Stard D Microwave Digestion Lab station with internal Temperature sensor and 260 terminal teach screen With HPR1000/10S High Pressure Segmented Rotar (Application Note HPR-EN-33).

Microwave Program 2 Steps (1) 15.00 Min (temperature 200⁰C) (2) 15.00 Min, (temperature 200⁰C).

After Finish left vessels 20 min until reach the room temperature, then the digested portion was diluted to a final volume of 50 ml using deionized water¹⁹. Pb and Cd Analyzed without Further Treatment. Fish samples were washed with deionized water, sealed in polyethylene bags and kept in a freezer at 20°C until chemical analysis²⁰.

Fish tissues were dried in oven at (80°C) until sample is at a constant weight. About 0.500 g of dry tissue sample (muscles, liver or gills) was accurately weighed and digested with 7ml of concentrated nitric acid (HNO₃ 65%) and 1ml of hydrogen peroxide (H₂O₂ 30%). Milestone Stard D Microwave Digestion Lab station with internal Temperature sensor and 260 terminal teach screen with HPR1000/10S High Pressure Segmented Rotar (Application Note HPR-FO-07) and AOAC Official Method 999.10²¹ and AOAC Official Method 974.14²² . Microwave Program 2 Steps (1)15.00 Min (temperature 200) (2)15.00 Min (temperature 200). After Finish left vessels 20 min until reach the room temperature, then the digested portion was diluted to a final volume of 50 ml using deionized water , before proceeding Method 3052²³. Pb Analyzed without further treatment, Cd diluted with Factor 2. The certified reference material DORM-2 analyzed for Pb and Cd content. Graphite furnace atomic absorption spectrometry (Model 220 GF), U.S.A Made, were used for analysis of Cd and Pb in fish tissue samples, Perfect for AOAC Official Method 999.10²¹. Wavelength, energy, lamp and burner alignment and slit width were optimized for Pb and Cd analysis (Table 1). For sea water and fish tissue samples five standard solutions were made (Table 2). Wavelength, energy, lamp and burner alignment and slit width were optimized for Pb and Cd analysis (Table 3). For sediments samples standard solutions were made (Table 4).

Statistics

All heavy metals data (lead and cadmium) were analyzed and tested for differences between group means of stations and seasons for significance

(P≤0.05)using the analysis of variance one way ANOVA and two ways ANOVA technique. Also, group means of environmental factors were analyzed by one way ANOVA technique. All statistical analysis was performed using the Origin 9 and SPSS software packages, version 17.0 .

RESULTS

Heavy metals in filtered water surface- The highest concentration of Pb in filtered water surface of Aden was 0.055 mg/L on winter 2011 and the lowest concentration was 0.045 mg/L on summer 2012. The highest concentration of Cd in filtered water surface of Aden was 0.010 mg/L on summer 2012 and the lowest concentration was 0.006 mg/L on winter 2011(Table 5).

Heavy metal in surface sediment- The highest concentration of Pb in sediments of Aden was 35.104 µg/g (dry wt.) on summer 2012 and the lowest concentration was 33.507 µg/g (dry wt.) on winter 2011 . The highest concentration of Cd in sediments of Aden was 2.111 µg/g (dry wt.) on summer 2012 and the lowest concentration was 1.775 µg/g (dry wt.) on winter 2011(Table 5).

Heavy metals in muscles fish- the highest concentration of Pb in the muscles of fish was 0.075 µg/g (dry wt.) on winter 2013 and the lowest concentration was 0.059 µg/g (dry wt.) on winter 2011. Also, the highest concentration of Cd in the muscles of fish was 0.052µg/g (dry wt.) on Summer 2012, whereas the lowest concentration was 0.024 µg/g (dry wt.) on winter 2011 (Table 5). However, the highest concentration of Pb in the muscles of fish was 0.116 µg/g (dry wt.) on large *E. areolatus* and the lowest concentration was 0.020 µg/g (dry wt.) on small *S. jello*. Also, the highest concentration of Cd in the muscles of fish was 0.073µg/g (dry wt.) On large *L. mahsena*, whereas the lowest concentration was 0.018 µg/g (dry wt.) on small *S. jello* (Table 6).

Heavy metals in liver- the highest concentration of Pb in the Liver of fish was 0.150 µg/g (dry wt.) on summer and the lowest concentration was 0.110 µg/g (dry wt.) on winter 2011 ; on the other hand, the highest concentration of Cd in the Liver of fish was 0.130 µg/g (dry wt.) on winter 2013, whereas the lowest concentration was 0.057 µg/g (dry wt.) on winter 2011 (Table 5). The highest concentration of Pb in the Liver of fish was 0.267µg/g (dry wt.) on large *L. mahsena* and the lowest concentration was 0.038 µg/g (dry wt.) on small *S. jello*. Also, the highest concentration of Cd in the Liver of fish was 0.172 µg/g (dry wt.) On Large *T. tonggol*, whereas the lowest concentration was 0.028µg/g (dry wt.) on small *S. jello* (Table 6).

Heavy metals in gill- The highest concentration of Pb in the Gill of fish was 0.290 µg/g (dry wt.) on Summer and the lowest concentration was 0.212 µg/g (dry wt.) on winter 2013. Also, the highest concentration of Cd in the Gill of fish was 0.348 µg/g (dry wt.) on summer, whereas the lowest concentration was 0.092 µg/g (dry wt.) on winter 2011(Table: 5). the highest concentration of Pb in the Gill of fish was 0.727 µg/g (dry wt.) on small *L. mahsena* and the

lowest concentration was 0.047 $\mu\text{g/g}$ (dry wt.) on small *S. jello*; but, the highest concentration of Cd in the gill of fish was 0.609 $\mu\text{g/g}$ (dry wt.) on small *E. areolatus*, whereas the lowest concentration was 0.033 $\mu\text{g/g}$ (dry wt.) on small *S. jello* (Table 6).

DISCUSSION

The heavy metals Pb and Cd were noticeably abundant in samples collected in the summer. The abundance of metals in filtered surface water samples is interpreted to be due to amount of draining sewage on summer were higher compared with winter and also due to high water temperature on summer season.

The heavy metals Pb and Cd were noticeably abundant in samples collected from sites located near the traffic ways and from areas near sewage discharge.

However, our results are in a good agreement with those found by²⁴ showed that the concentration of Pb was 0.034 ± 0.002 mg/L and Cd was 0.012 ± 0.001 mg/L in water from the Kolleru Lake, India, on summer. Besides²⁵, pointed out that the concentration of Pb was 0.03-0.07 mg/L, which is below the permissible limit of 0.1 mg/L set for inland surface water, in the water samples collected from sea water in Międzyzdroje, Baltic coast, Poland

Heavy Metals in Surface Sediments, The highest concentration Pb and Cd were in the surface sediments of Aden coast was obtained on summer. This result may be explained by the fact that during summer that region faced with decreasing internal currents and water supplies that might have caused increase of heavy metals comparison with winter season, wastewater discharge from the city. The obtained results for Cadmium and their interpretation are comparable to the concentration of Cd was 0.20- 5.80 $\mu\text{g/g}$ (dry wt.)²⁶ in summer from Red Sea coast, Al Hodeidah, Yemen, and a good agreement with those found by²⁷ pointed out that the concentration of Pb was 61.620 $\mu\text{g/g}$ (dry wt.) in northern Delta Lakes, Egypt. **Heavy metals in Muscles Fish,** The highest concentration of Cd in muscles of fish species studied was on summer. The present high concentration of Cd in summer may be attributed to various factors as sewage, increase in the water temperature during summer increases heavy metals uptake in the fishes as compared to the winter season, higher metabolic rate in the fishes. The highest concentration of Pb and Cd in muscles of fish of *Epinephelus areolatus* and *Lethrinus mahsena*. The present high concentration of Pb and Cd in *E. areolatus* and *L. mahsena* have been usually attributed to their habitat and feeding behavior. *E. areolatus* and *L. mahsena* tend to be near the sediment region.

The present high concentration of Pb and Cd in Aden may be due to the anthropogenic activities, feeding behavior of the fish species, fat content, the in their diet uptake. Some of these values are very small compared with the obtained readings in the current study, however, not be surprising because the marine environment of the ocean is generally less polluting for the sea that are surrounded by industrialized countries, or that are over a lot of ships, such as the Gulf of Aden line. The mean concentrations of Pb and Cd were

considered low and under the recommended safe levels set by [FAO/WHO²⁸ and Yemen standard²⁹.

However, our results are in a good agreement with those found³⁰, showed that the concentration of Pb was 0.100 ± 0.09 $\mu\text{g/g}$ (dry wt.) in muscles of *Epinephelus areolatus* from Tuticorin, India. Besides,³¹ pointed out that the concentration of Pb was 0.20 ± 0.02 $\mu\text{g/g}$ (dry wt.), in *L. mahsena* from Red Sea, Egypt³² pointed out that the concentration of Pb was 0.118- 0.193 $\mu\text{g/g}$ (wet wt.) and Cd was 0.013- 0.023 $\mu\text{g/g}$ (wet wt.), in tuna fish Hadhramout Coast Yemen; ³³ pointed out that the concentration of Pb was 0.12 ± 0.21 $\mu\text{g/g}$ (dry wt.), in *L. mahsena* from Red Sea of Al-Cornish Hodeidah Yemen; pointed out³⁴ that the concentration of Cd was 0.09 ± 0.010 $\mu\text{g/g}$ (wet wt.), in *T. albacares* from Aden Yemen;³⁵ pointed out that the concentration of Pb was 0.10 - 2.10 $\mu\text{g/g}$ (dry wt.), Cd was 0.06-1.06 $\mu\text{g/g}$ in *L. mahsena* from Jeddah Coast, Saudi Arabia.

Heavy metals in Liver Fish, The highest concentration of Pb and Cd in Liver of fish, were observed to be more concentrated in the larger sizes of fish. The present high concentration of Pb and Cd in the larger sizes of fish may be attributed to various factors as large fish that prey upon smaller fish can accumulate more of the chemical in their bodies. It is better to eat the smaller fish within the same species, the strong affinity of metallothionein protein with these elements. This is usually more pronounced in bigger fishes. The highest concentration of Cd in Liver of fish species studied was on summer.

The present high concentration of Cd in summer may be attributed to various factors as differences in local pollution, industrial wastes, bioavailability of metals (variations among physiochemical factors) and fish metabolism (growth cycle, reproduction and feeding). physiological changes, minor role of annual cycles of pH and metal concentration in the water and metal level of the diet in the seasonal pattern of metal concentration in liver.

The highest concentration of Pb in Liver of *L. mahsena*, but, the highest concentration of Cd in the Liver of *E. areolatus*. The present high concentration of Pb and Cd in *L. mahsena* and *E. areolatus* may be attributed to various factors, demersal or bottom-dwelling species, food preferences, organism mobility or other attributes of behavior with respect to the environment, strong binding with cystine residues of MT, lipid content in the tissue and excretion percentage of these toxic metals from their body, increased metabolic rate, water temperature. However, our results are in a good agreement with those found by³⁶ pointed out that the concentration of Cd was 0.6 $\mu\text{g/g}$ (dry wt.) in liver of *Epinephelus fasciatus* from Jordan; and³⁷ pointed out that the concentration of Cd was 0.25 ± 0.04 $\mu\text{g/g}$ (dry wt.) in liver of *E. areolatus* from Hong Kong, China. The highest concentration of Cd was found in Gill of fish on summer. These observations probably as indicated in the case of liver tissue which mentioned above, beside that in water, gills are the main surface during exchange of ions metals. The highest concentration of Pb in Gill of fish of *L. mahsena*, but, the highest concentration of Cd and As

in the Gill of fish of *E. areolatus*; on the other hand, the highest concentration of Hg in the Gill of fish of *T. tonggol*.

The present high concentration of Pb in *L. mahsena* may be attributed to various factors, the metal complexation with the mucus that is impossible to remove completely from the lamellae before analysis, the similarity of lead and calcium in their deposition and mobilization from the gill, the result of a water contamination caused by environmental pollution. However, our results are in a good agreement with those found by³⁸ showed that the concentration of Cd was 0.12-0.35 µg/g (dry wt.) in Gill of *Mystusseenghala* and *Wallagoattu* from Pakistan³⁹, in summer; and showed that the concentration of Cd was 0.300±0.01 µg/g (dry wt.) in Gill of *Scardinius erythrophthalmus* from the Topolnitsa reservoir (Bulgaria), on summer; and another study⁴⁰ showed that the concentration of Cd was 0.34 ± 0.20 µg/g (wet wt.) in Gill of *Cyprinus carpio* from Sikkak dam at Ainyoucef (Wilaya of Tlemcen) Algeria, in summer.

Similar results are found in high Pb concentrations in gills were recorded by^{36, 41, 42, 43}.

However, our results are in a good agreement with those found by³³ showed that the concentration of Pb was 1.20±0.95-1.70±0.90 µg/g (wet wt.) in Gill of *Lethrinus lentjan* and Cd was 0.25±0.07 - 0.54±0.10 µg/g (wet wt.) in Gill of *Epinephelus sexfasciatus* During winter from Red Sea, AL-Hudaydah, Yemen; and³¹ showed that the concentration of Pb was 0.23±0.02 µg/g (dry wt.) in Gill of *L. mahsena* and Cd was 0.54±0.07 µg/g (dry wt.) in Gill of *Epinephelus spp* from Red Sea, Egypt; and⁴⁴ showed that the concentration of Cd was 0.117±0.087 µg/g (dry wt.) in Gill of *Epinephelus coioides* from northern of Persian Gulf.

The present high concentration of Cd in *E. areolatus*; suggesting that their gills could accumulate trace metals from the environment. *E. areolatus* are both omnivorous feeders (the former being a bottom feeder), metabolism, biological and ecological factors such as feeding habits and habitat, have a close relationship with sediment. A possible explanation for this might be to electricity generating stations cooling waters and effluents, sewage disposals and storm waters, Another possible explanation for this is that Scrap-iron store at Labour Island is the most likely source of Pb and Cd in the Seawater, Oil Harbour and municipal sewages are expected sources to be responsible for elevated tissue Cd concentrations in Seawater from Sahel Abyen and Sira Island. This result may be explained by the fact that Aden area was the highest among the other stations. It is assumed that the presence of the area near the strait of Bab Al-Mandab, where the water coming from both the Red Sea and the Indian Ocean mixes and consequently changes its nature, especially during the seasonal monsoons, is responsible mainly for elevated Gill tissue Cd concentrations in the fish inhabiting the Aden area. However, some sewage outflow can be additional anthropogenic sources of Cd in the area. Apparently, Cd contamination gradients exist in the Gulf of Aden waters.

CONCLUSION

In conclusion, heavy metals accumulate in various tissues of fish with different amount. Overall, accumulation of metals in muscle was lower than liver and gills. The results present new information on the distribution of these metals in liver, gills and muscle of *Lethrinus mahsena*, *Thunnus tonggol*, *Sphyræna jello* and *Epinephelus areolatus*. Generally, this research showed that concentrations of heavy metal in the Aden Coast, Gulf of Aden, Yemen were so far significantly lower than effects range low (ERL) and lower than the maximum permissible concentration for various countries. According to the fish samples analyses, concentrations of heavy metal in fish species tissues were well within the limits set by the (FAO/WHO (2004) and Standard Specification for Yemen, 2006). Recommendations and showed that the fish from investigated region are safety for consumers.

ACKNOWLEDGEMENT

Our great thanks are for those enabling us to accomplish this work, the patients. Without them, this work could not be carried out and we would like to acknowledge with much appreciation to all people and technical in the Faculty of Environmental Sciences and Marine Biology Laboratories, Hadramout University, laboratory at the Royal Scientific Society in Jordan (RSS), laboratory at the Dr. Babasaheb Ambedkar Marathwada University in India (BAMU), University and all technical in Chemical Laboratories in University of Science and Technology- Aden Branch.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. Abdel-Baki AS, Dkhil MA, Al-Quraishy, S. Bioaccumulation of some heavy metals in tilapia fish relevant to their concentration in water and sediment of Wadi Hanifah, Saudi Arabia. African J Biotech. 2011; 10(1): 2541-2547.
2. Belin S, Sany T, Salleh A. Heavy metal contamination in water and sediment of the Port Klang coastal area, Selangor, Malaysia. 2013; (1):2013-2025
3. Wu Y, Chen J. Investigating the effects of point source and nonpoint source pollution on the water quality of the East River (Dongjiang) in South China. Ecological Indicators. 2013; 32(1):294-304.
4. Varol M, Şen B. Assessment of nutrient and heavy metal contamination in surface water and sediments of the upper Tigris River, Turkey. CATENA. 2012; 92(1):1-10.
5. Wang C, Niu Z, Li Y, Sun J, Wang F. Study on heavy metal concentrations in river sediment through the total amount evaluation method. J Zhejiang Univ.-Sci. A Appl. Phys. and Eng. 2011; 12(5):399-404
6. Saeed S M, Shaker SF. Impact of cage-fish culture in the river Nile on physico-chemical characteristics of water, metals accumulation, histological and some biochemical parameters in fish. Abbassa Int. J. Aqua. 2008; (1A): 179-202.
7. Dean TA, Bodkin JL, Fukuyama AK, Jewett SC, Monson, DH, O'Clair CE, Van Blaricom GR. Food limitation and the recovery of sea otters following the *exxon valdez* oil spill. Marine Ecology Progress Series. 2002; 241(1):155-270.
8. Kwon YT, Lee CW. Sediment metal speciation for the ecological risk assessment. Anal. Sci. 2001; 7(1):1015-1017
9. Olaifa F E, Olaifa AK, Adelaja AA, Owolabi, AG. Heavy metal concentration of *Clarias gariepinus* from a lake and

- fish from in Ibadan, Nigeria. Afr J Biomed. Res, 2004; 7(1):145-148
10. Burger J, Gaines KF, Boring CS, Stephens W L, Snodgrass J, Dixon C, McMahon M, Shukla S, Shukla T, Gochfeld M. Metals levels in fish from the Savannah River: Potential hazards to fish and other receptors. Environ. Res. 2002; 89(1):95-97
 11. Clearwater SJ, Baskin SJ, Wood CM, Mac Donald. Gastrointestinal uptake and distribution of copper in rainbow trout. J Exp Biol. 2000; 203(1):2455-2466
 12. Van den Broek JI, Gledhill KS, Morgan DG. Heavy metal concentration in the Mosquito fish *Gambusia holbrooki* in the manly Lagoon Gatchment. In: UTS, fresh water ecological report 2002 department of environmental Science, University of Technology, Sydney. 2002; (1):1-25
 13. Schulz UH, Martins-Junior, H. Astyanafaseiatus as bio indicator of water pollution of Rio Dos Sinos, Rs, Brazil. Braz J Biol. 2001; 61(4):615-622
 14. Oforka NC, Osuiji LC, Onwuachu UI. Estimation of Dietary intake of cadmium, lead, manganese, zinc and nickel due to consumption of chicken meat by inhabitants of Port-Harcourt Metropolis, Nigeria. Archives of Applied Science Research. 2012; 4(1):675-684.
 15. Mason LH, Harp JP, Han DY. Pb Neurotoxicity: neuropsychological effects of lead toxicity. Biomed. Res. Int.; Vol. 2014
 16. Oronsaye JAO, Wangboje OM, Oguzie FA. Trace metals in some benthic fishes of the Ikpobariver dam, Benin City, Nigeria; African J Biotechnology. 2010; 9(1):8860-8864.
 17. Dural M, Goksu LM, Ozak AA. Investigation of heavy metal levels in economically important fish species captured from the Tuzla lagoon. Food Chemistry. 2007; 102(1):415-421.
 18. US EPA. Method 200.13: Determination of Trace Elements in Marine Waters by Off-Line Chelation Pre concentration with Graphite Furnace Atomic Absorption. Washington, D.C.: U.S. Environmental Protection Agency, 1992.
 19. US EPA. Method 3051A Microwave assisted acid digestion of sediments, sludge, soils, and oils. Washington, D.C.: U.S. Environmental Protection Agency, 2007.
 20. US EPA. Environmental Protection Agency Chemical Contaminant Data for Use in Fish Advisories Vol. 1 Fish Sampling and Analysis Third Edition. Washington, DC: Office of Science and Technology (2000).
 21. AOAC. Official Method 999.10. Lead, cadmium, zinc, copper and Iron in foods, atomic absorption spectrophotometry after microwave digestion, first action 1999. NMKL-AOAC method, 2005.
 22. AOAC, Association of Official Analytical Chemists: Official Method 974.14. Mercury in Fish, first action 1974. NMKL-AOAC method, Washington, DC, USA, 18th edition, Revision 3, 2010.
 23. US EPA. Method 3052 microwave assisted acid digestion of siliceous and organically based matrices. Washington, D.C.: U.S. Environmental Protection Agency, 1996.
 24. Mastan SA. Heavy metals concentration in various tissues of two freshwater fishes, *Labeo rohita* and *Channa striatus*. African J Environmental Science and Technology. 2014; 8(2):166-170.
 25. Daniszewski P. Determination of metals in sea water of the Baltic Sea in Międzyzdroje, International Letters of Chemistry, Physics and Astronomy. 2013; 18(1):13-22
 26. Heba HM, AL-Edresi AM, Al-Saad H, Abdolmoneim. A. Background Levels of Heavy Metals in Dissolved, Particulate Phases of Water and Sediment of Al-Hodeidah Red Sea Coast of Yemen. J. King Abdulaziz Univ. Mar. Sci. 2004; 15(1):53-71.
 27. Saeed SM, Shaker SF. Impact of cage-fish culture in the river Nile on physico-chemical characteristics of water, metals accumulation, histological and some biochemical parameters in fish. Abbassa Int J Aqua. 2008; 1(A): 179-202.
 28. FAO/ WHO. Summary of Evaluations Performed by the Joint FAO/WHO Expert Committee on Food Additives (JECFA 1956-2003), ILSI Press International Life Sciences Institute. 2004.
 29. Yemen Standardization, Metrology and Quality Control. Fish and fish products, fresh fish. The Republic of Yemen. 2006; 1577(1):1-9.
 30. Pugalandhi T, Uma Maheswari, G. Concentration of lead and cadmium in some edible fishes from Tuticorin. J Mar Biol Ass. India. 2007; 49(2): 254-256
 31. Al-Ghanim KA, Abdelatty M, Abdelfattah L, Mahboob S. Studied the differential uptake of heavy metals by gill, muscles and liver of four selected fish species from Red Sea. Pakistan J Zool. 2015; 47(4):1031-1036
 32. Algahri MA, Bamoteref S Kh, Saeedan AM. Lead, Mercury and Cadmium in Tuna Fish Caught at the coast of Hadramout – Yemen. Hadramout University J Natural Applied Sciences. 2011; 8(2):225.
 33. Al-Zubaidy AB, Majam MT, Faqeh EA. Investigation of heavy metals contamination and parasites of edible marine seafood Yemeni, Coastal Waters. J Purity, Utility Reaction and Environment. 2014; 3(6):128-145
 34. Larouci Mohammed Ben Yehya. A Study on Contamination Levels with Heavy Metals in some Fish and Shellfish Species in Yemeni Regional Sea. M.Sc. thesis, Sana'a University. 2006.
 35. Ali AA, Elazein EM, Alian MA. Investigation of heavy metals pollution in water, sediment and fish at red sea–Jeddah Coast- KSA at two different locations. J Applied Environmental and Biological Sciences. 2011;1(12):630-637
 36. Abu Hilal AH, Ismail NS. Heavy Metals in Eleven Common Species of Fish from the Gulf of Aqaba, Red Sea. Jordan J Biological Sciences. 2008; 1(1):13 – 18.
 37. Wong CK, Wong PPK, Chu LM. Heavy metal concentrations in marine fishes collected from fish culture sites in Hong Kong. Archi Envi Contamin Toxicol. 2001; 40(1):60-69.
 38. Inayat I, Batool AI, Rehman MF, Ali NHNA, Jabeen SH. Seasonal bioaccumulation of heavy metals in the right and leftgills of edible fishes. World J Fish Marine Sci. 2014; 6 (2):195-200
 39. Yancheva V, Stoyanova S, Velcheva I, Petrova S, Georgieva, E. Metal bioaccumulation in common carp and rudd from the Topolnitsa reservoir, Bulgaria, Arh Hig Rada Toksikol. 2014;65(1):57-66
 40. Dali Youcef D, Nacéra, Mesli Lotfi. Seasonal Variations Of Heavy Metals In Common Carp (*Cyprinus Carpio* L., 1758) Collected From Sikkak Dam Of Tlemcen (Algeria). J Engineering Research and Applications. 2014; 4(1):01-08
 41. Kargin F. Metal concentrations in tissues of the freshwater fish *Capoeta barroisi* from the Seyhan River (Turkey). Bull. Environ. Contam. Toxicol. 1998, 60(1):822-828.
 42. Avenant-Oldewage A, Marx HM. Bioaccumulation of chromium, copper and iron in the organs and tissues of *Clarias gariepinus* in the Olifants River, Kruger National Park, Water SA-Pretoria. 2000;26(4):569-580
 43. Qadir A, Malik R. Heavy metals in eight edible fish species from two polluted tributaries (Aik and Palkhu) of the River Chenab. Pakistan, Biol Trace Elem Res. 2011; 143(1):1524-1540.
 44. Khoshnood Z, Khoshnood R, Mokhlesi. A.; Ehsanpour, M.; Afkhami, M. and khazaali, A. Determination of Cd, Pb, Hg, Cu, Fe, Mn, Al, As, Ni and Zn in important commercial fish species in northern of Persian Gulf. J Cell Animal Biol. 2012; 6(1):1-9

Table 1: Operational parameters for Pb and Cd analysis by GFAAS in surface seawater and Fish

Element	Pb	Cd
Sample	Sw+F	Sw+F
Method	GFAAS	GFAAS
Wavelength (nm)	283.3	228.8
T Drying ^a , °C	110-130	110-130
T Pyrolysis, °C	1000	900
T Atomization, °C	1900	1200
T Clean out, °C	2500	2500
T Ramp, °C	1400	1300
Matrix modifier	5 µL NH ₄ H ₂ PO ₄ +Mg(NO) ₂	5 µL NH ₄ H ₂ PO ₄ +Mg(NO) ₂
Slit width (nm)	0.8	1.2
Argon flow (ml min ⁻¹)	250	250
Injection volume (µL)	20	20
Instrumental detection	0.02	0.005

Sw Sea water, F fish, GFAAS graphite furnace atomic absorption spectrometry

Table 2: Standard Concentration for Pb and Cd Analyzed in Surface Seawater and Fish

Element	Pb	Cd
Sample	Seawater+ Fish	Seawater+ Fish
Method	GFAAS	GFAAS
Standard Solutions (µg/L)	0.5, 1.0, 1.5, 2.0, 2.5	0.125, 0.25, 0.50, 0.75, 1.0

Table 3: Operational parameters for Pb and Cd Analysis by FAAS in Surface Sediment

Element	Pb	Cd
Sample	Sediment	Sediment
Method	Flam	Flam
Wavelength (nm)	283.3	228.8
Flam mode	C ₂ H ₂ /Air	C ₂ H ₂ /Air
Slit width (nm)	0.7	0.7
Lamp current (mA)	2	2
Energy (eV)	2.874	3.214
Instrumental detection limit (mg/L)	0.08	0.01

Table 4: Standard Concentration for Metals Analyzed in Surface Sediment

Element	Pb	Cd
Sample	Sediment	Sediment
Method	flam	flam
Standard Solutions (mg/L)	1.0, 2.0, 4.0, 8.0	0.05, 0.1, 0.15, 0.20, 0.25

Table 5: The mean of concentration (mg/L) for lead and cadmium during the seasons in water, sediments and fish organs (muscle, liver and gill) of Aden station, Yemen coast.

Samples	Metal ion	Seasons			Total mean±SD
		Winter 2011	Summer 2012	Winter 2013	
Filtered surface water	Pb	0.055±0.004	0.045±0.007	0.051±0.005	0.050±0.005
	Cd	0.006±0.002	0.010±0.003	0.009±0.001	0.008±0.002
Sediments	Pb	33.507±1.132	35.104±0.416	34.688±1.285	34.433±0.828
	Cd	1.775±0.201	2.111±0.839	2.045±0.310	1.977±0.238
Muscles	Pb	0.059±0.028	0.071±0.032	0.075±0.035	0.068±0.031
	Cd	0.024±0.010	0.052±0.021	0.041±0.019	0.039±0.015
Liver	Pb	0.110±0.042	0.150±0.096	0.144±0.086	0.135±0.072
	Cd	0.057±0.026	0.126±0.093	0.130±0.070	0.104±0.058
Gill	Pb	0.243±0.225	0.290±0.366	0.212±0.219	0.248±0.267
	Cd	0.092±0.075	0.348±0.354	0.310±0.296	0.250±0.228

Table 6: Lead and Cadmium concentrations (µg/g) in muscles, Liver and Gill samples

Species	Size	Muscles		Liver		Gill	
		Pb (µg/g)	Cd (µg/g)	Pb (µg/g)	Cd (µg/g)	Pb (µg/g)	Cd (µg/g)
<i>Lethrinus mahsena</i>	Large	0.09±0.012	0.073±0.021	0.267±0.076	0.107±0.013	0.665±0.138	0.095±0.016
	Medium	0.078±0.011	0.045±0.013	0.23±0.088	0.095±0.015	0.672±0.164	0.08±0.009
	Small	0.071±0.018	0.034±0.006	0.22±0.082	0.087±0.012	0.727±0.257	0.058±0.013
<i>Thunnus tonggol</i>	Large	0.083±0.009	0.043±0.02	0.137±0.012	0.172±0.069	0.133±0.054	0.518±0.18
	Medium	0.079±0.011	0.031 ± 0.01	0.123±0.01	0.099±0.048	0.123 ±0.049	0.34±0.177
	Small	0.058±0.013	0.025±0.009	0.103±0.011	0.074±0.038	0.103 ±0.032	0.16±0.07
<i>Sphyraena jello</i>	Large	0.028±0.006	0.03± 0.015	0.068±0.01	0.038±0.015	0.079 ±0.013	0.051±0.019
	Medium	0.021±0.002	0.024±0.012	0.044±0.008	0.032±0.016	0.058 ±0.005	0.038±0.006
	Small	0.02 ±0.002	0.018 ± 0.005	0.038±0.007	0.028±0.01	0.047 ±0.007	0.033±0.003
<i>Epinephelus areolatus</i>	Large	0.116±0.007	0.055 ± 0.025	0.141±0.01	0.164±0.084	0.132 ±0.008	0.512±0.369
	Medium	0.093±0.026	0.05 ± 0.026	0.129±0.006	0.145±0.076	0.125±0.009	0.507±0.388
	Small	0.083±0.026	0.037 ± 0.021	0.122±0.006	0.209±0.139	0.111± 0.007	0.609±0.469