Available online on 15.05.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal
 Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial Share Alike 4.0 License which permits unrestricted non commercial use,

provided the original work is properly cited

Volume 5, Issue 2, 2020

Open Access

REVIEW ARTICLE

CRITICAL REVIEW ON *BOMBAX CEIBA*, *ALOE VERA* AND *XIMENIA AMERICANA*

Rohit Shankar Mane , Ankala Basappa Vedamurthy

Department of Biotechnology and Microbiology, Karnatak University, Dharwad, Karnataka, 580003, India.

ABSTRACT

Therapeutic plants are extensively increased in scenery with their efficient therapeutic properties. India has the greater natural ecosystem from past two decades, near has been a marvelous augment in the applications of natural medicines; though there is still a noteworthy deficiency in the research of medicinal plants and it has the great resources of medicinal plants which are noteworthy to human beings in many ways. The medicinal plants such as *Bombax ceiba*, *Aloe vera* and *Ximenia Americana* are well known to cure several diseases therefore used in different Ayurvedic preparation somehow they are lagging behind in the list of medicinal plants for their applications in drug preparations. These plants has great ethnopharmaceutical and pharmacological properties with therapeutic applications therefore present review gives different aspects of *Bombax ceiba*, *Aloe vera* and *Ximenia americana* and may help to create awareness in society.

Key words: *Aloe vera*, *Bombax ceiba*, ecology, ethnobotany, pharmacology, *Ximenia americana*.

Article Info: Received 2 February 2020; Revised 6 April; Accepted 7 May, Available online 15 May 2020

Cite this article-

Rohit Shankar Mane, Ankala Basappa Vedamurthy. Critical review on *Bombax ceiba*, *Aloe vera* and *Ximenia americana*. Universal Journal of Pharmaceutical Research 2020; 5(2):65-67.

DOI: <https://doi.org/10.22270/ujpr.v5i2.392>

Address for Correspondence:

Dr. Ankala Basappa Vedamurthy, Department of Biotechnology and Microbiology, Karnatak University, Dharwad, Karnataka, 580003, India. Tel: +6281524045514, E-mail: vedamurthybt@gmail.com

INTRODUCTION

The British biologist Norman Myers coined the term "biodiversity hotspot" in 1988. He described them as a biogeographic area by characterizing their brilliant steps of plant endemism and grave levels of habitat loss¹. Further, Conservation International (CI) joined 'Myers hotspots criteria' in 1989-1996, and they made one organization for the reassessment of the hotspots concept and in 2005 they have in print an efficient posh Hotspots Revisited: Earth's Biologically Richest and Most Endangered Earthly Ecoregions². According to the CI report, there is the total of thirty-five biodiversity hotspots in the world, amongst them, four are in India. Mainly these three biodiversity hotspots are situated in the Himalaya, Indo-Burma, Sundaland, and the Western Ghats³. A Total of 17000-18000 flowering plant species, 8000 medicinal plants, are documented in folk and Ayurveda, Unani, Siddha, and Homoeopathy⁴. Herbs are the immense reserve base for the conventional medication & herbal industry and also it admits livelihood and health safety to a great section of Indian population. India has the greater natural ecosystem from past two decades, near has been a marvelous augment in the applications of natural medicines; though there is still a noteworthy deficiency in the research of medicinal plants and it has the great resources of medicinal plants which are noteworthy to

human beings in many ways⁵. According to the WHO, herbs are used as the best resource to get a diversity of bioactive compounds in the development of different drugs, those are effective as an antimicrobial, anti-tuberculosis, antioxidant, anticancerous, anti-inflammatory, antidiabetic, anthelmintic, hepatoprotective activity, larvicidal activity⁶⁻⁹. By taking the above background in the present review total of three medicinal plants revealed with their critical information.

In the present study a total of three medicinal plants were under study and their information is as follows.

BOMBAX CEIBA LINN

Bombax ceiba diversified under the family Bombacaceae. It is a bulky, deciduous, elegant plant inhabiting Africa, Australia, temperate and tropical Asia¹. It has been applied in traditional medicine for its versatility in nature against pathogens²⁻⁴. These plant roots and flowers were used to treat different ailments. The plant has different vernacular names such as Semal, Silk-cotton tree (English), Sembhal (Urdu), Semar (Hindi), Buruga (Telugu), Simalu (Assam), Moca (Sanskrit), Shimool (Bengali), Shemalo (Gujarati)⁴.

The plant botanical classification is as follows:

Kingdom: Plantae
Division: Magnoliophyta
Class: Magnoliopsida
Order: Malvales
Family: Bombacaceae
Genus: Bombax
Species: ceiba

It is extensively originated in moderate Asia, Africa, Australia and tropical Asia. In India, it can be originated at altitudes up to 1500m. It is usually seen within the arid and wet deciduous forests and also by rivers of peninsular India. It is a huge light-eater and rapidly rising tree. It grows finest on bottomless grimy loams, especially in valleys, within the parts that are receiving 50 to 460 cm⁵. The morphology is shown in Figure 1. The dissimilar pieces of *B. ceiba* similar to all parts are revealed to hold various curative behaviors in pharmlological surveys. In February, *B. ceiba* begins

plummeting every one of its foliage. It is time for blossoming and follows an amazing exhibit of great silky red flowers at the tips of naked branches. In May, white cottony strands, from opened fruits, float downward, settling on the ground, houses and whatever else is in their mode. It displays spiky twigs orderly in horizontal tiers which are rough, straight and spiny with buttress roots¹⁻³. The body parts of the plant contain different chemicals. Bark contains lupeol, saponins, tannins, gums and trihydroxyflavone-3-O-β-D-glucopyranosyl(1-4)-α-L-rhamnopyranoside, nhexacosanol, shamimicin; Seeds contain palmitic acid; roots contains lactone, Hemigossypol-6-methyl; flowers contains polysaccharides; Leaves contain a flavonols, C-glycoside shamimin and Xanthone. The plant has different medicinal activities such as anti-inflammatory, treating leprosy, muscular injury, wounds, asthma, birth control, sexual diseases¹⁻⁵.

Figure 1: Leaves, stem, roots and seeds of *Bombax ceiba*.

Figure 2: Leaves, pulp and roots of *Aloe vera*.

ALOE VERA

The plant comes under the Asphodelaceae family. It is a shrubby. It is perennial. It is xerophytic and succulent. It is pea green coloured plant. It mainly founds in the dry regions of the world. These dry regions are nothing but the regions from America, Asia, Africa and Europe. Inside India, the plant is found in Maharashtra, Gujarat, Andhra Pradesh, Tamil Nadu and Rajasthan⁶. It has leaves and these leaves are fleshy and triangular in nature. The leaves are jagged with edges. It consist yellow tubular flowers. It also consist fruits that contain frequent seeds⁷. The morphology is shown in Figure 2. Leaves are made up of three layers. Inner layer is made up of clear gel; middle layer is made up of latex while outer layer is made up of cells. The outer layer is thick in nature due to carbohydrates and proteins.

Taxonomy

Kingdom: Plantae   **Order:** Asparagales
Division: Spermatophyte   **Genus:** Aloe
Subdivision: Angiospermae
Class: Monocotyledoneae
Species: Barbadosensis Mill

Synonyms: Aloe, Musabbar, Kumari

The plant contains nataloins and barbaloins. These two aloins are important in the view of medicinal applications. Nataloins gives picric acids and oxalic acids in the presence of nitric acid. It will not give a red coloration after reaction. Barbaloins gives aloetic acid and chrysammic acid in the presence of nitric acid. Barbaloin yields yellow coloured prismatic crystals⁶⁻⁸. *Aloe vera* produces six antiseptic agents. These agents are active against all pathogens and nothing but cinnamonic acid, lupeol, phenols and sulphur, salicylic acid and urea nitrogen. It is highly active in the action of scrapes, burns, abrasions, cuts, rheumatoid arthritis, allergic reactions, rheumatic fever, ulcers, acid indigestion, and many more inflammatory conditions of the digestive system. It also shows β- sitosterol and it is also powerful anti-cholesterol agent. It helps to lower damaging cholesterol levels⁶⁻⁹.

XIMENIA AMERICANA

The *Ximenia americana* tree belongs to the Olacaceae family. It is mainly originated in Jabal, Marra and Radom which comes under Darfur. It is made up of

leaves, bark and fruits. They have numerous uses in local medicine¹⁰. The leaves and twigs extracts are used to cure fever, cold, as a mouth wash for tooth aches and as a laxative and an eye lotion¹¹. The leaves are used for headaches and poison antidote¹¹. The morphology is shown in Figure 3.

Figure 3: Leaves, stem, roots and seeds of *Ximenia americana*.

The plant contains big bulgy roots. These roots are used in the treatment of skin aches, headaches, and hemorrhoids, sexually transmitted diseases, sleeping thickness, leprosy, guinea worm and oedema. It is rich in fruits. Fruits are used in the treatment of constipation. It consist bark. The dried bark transformed into powder and used to treat skin ulcers. The plant is rich in hydrocyanic acid. In the bark of the plant contains approximately 17% oils is obtained. The heartwood and flowers also contain essential oils. Plant parts contain different constituents used in folk medicines¹³. It was reported that *X. americana* seed oil contains acids. The most important ingredients of the volatile oil were benzaldehyde, hydroxy benzyl cyanide and isophorone¹². Stem of the plant consist steroids, triterpenoids, oleanolic acid, 3-oxo-oleanolic acid and sesquiterpenoids which were proved by ethanolic and methanolic extract¹². A large number of sesquiterpene are large number of constituents of higher plants. Terpenoids and lipophilic aromatic compounds were involved in the liverworts biological activities. Steroids and diterpenes are the secondary bioactive compounds of the plant. Plant chemically made up of saponins, glycosides, flavonoids, tannins, phenolics, alkaloids, quinones and terpenoids. These all above secondary bioactive compounds employed in different biological activities such as, antimicrobial, antifungal, anticancer, antineoplastic, antitrypanosomal, antirheumatic, antioxidant, and analgesic, and molluscicide, pesticide, also having hepatic and hematological effects¹⁰⁻¹³.

CONCLUSION

Present review article aimed to provide ill-intentioned, invasive, deleterious selected plants information to the readers. The knowledge about these three medicinal plants was insufficient in the literature; however this article will help to reveal more about the same plants. The ethanopharmaceutical and pharmacological studies revealed different aspects of *Bombax ceiba*, *Aloe vera* and *Ximenia americana* therefore present study may helpful to create awareness in society about the plants.

And the article will help to clear different myths about plants and it will help to upgrade the same plants in the pharmaceutically vital list of medicinal plants. Further, the same plants may helpful in the drug discovery and drug development therefore one can use present data.

AUTHOR'S CONTRIBUTION

All authors have worked equally for this work.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to Karnatak University to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

1. Ravi V, Patel SS, Verma K, Dutta D, Saleem TS. Hepatoprotective activity of *Bombax ceiba* Linn against Isoniazid and Rifampicin induced toxicity in experimental rats. *Int J Appl Res Nat Prod* 2010; 3(3): 19-26.
2. Sarita K, Greindra G, Shailsh G. Pharmacognostic and pharmacological profile of *Bombax ceiba*. *Asian J Pharm Educ Res* 2017; 6(3): 16-27.
3. Pankaj HC, Somshekhar SK. *Bombax ceiba* Linn.: Pharmacognosy, ethnobotany and phyto-pharmacology. *Pharmacog Commun* 2012; 2(3): 2-9. <http://dx.doi.org/10.5530/pc.2012.3.2>
4. Verma R, Devre K, Gangrade T, Gore S, Gour S. A pharmacognostic and pharmacological overview on *Bombax ceiba*. *Sch Acad J Pharm* 2014; 3(2): 100-107.
5. Kumar A, Lingadurai S, Jain A, Barman NR. *Erythrina Variegata* Linn: A review on morphology, phytochemistry and pharmacological aspects. *Pharmacogn Rev* 2010; 4(8): 147-152. <http://dx.doi.org/10.4103/0973-7847.70908>
6. Chithra P, Sajithlal G, Chandrakasan G. Influence of *Aloe vera* on the glycosaminoglycans in the matrix of healing dermal wounds in rats. *J Ethnopharmacol* 2018; 59: 179-86. [https://doi.org/10.1016/S0378-8741\(97\)00112-8](https://doi.org/10.1016/S0378-8741(97)00112-8)
7. Roberts DB, Travis EL. Acemannan containing wound dressing gel reduces radiation induced skin reactions in C3H mice. *Int J Radiat Oncol Biol Phys* 2019; 32: 1047-52. [https://doi.org/10.1016/0360-3016\(94\)00467-Y](https://doi.org/10.1016/0360-3016(94)00467-Y)
8. Sato Y, Ohta S, Shinoda M. Studies on chemical protectors against radiation XXXI: Protective effects of *Aloe arborescens* on skin injury induced by X-irradiation. *Yakugaku Zasshi* 2016; 110: 876-84. https://doi.org/10.1248/yakushi1947.110.11_876
9. Byeon S, Pelley R, Ullrich SE, Waller TA, Bucana CD, Strickland FM. *Aloe barbadensis* extracts reduces the production of interleukin after exposure to ultraviolet radiation. *J Invest Dermatol* 1988; 110: 811-7. <https://doi.org/10.1046/j.1523-1747.1998.00181.x>
10. Mahato SB, Nandy AK, Roy G. Triterpenoids. *Phytochemistry* 1992; 7(10): 180-187.
11. Maikai VA, Maikai BV, Kobo PI. Antimicrobial properties of stem bark extracts of *Ximenia americana*. *J Agri Sci* 2009; 1(2): 30-34. <https://doi.org/10.5539/jas.v1n2p30>
12. Fatope MO, Adoum OA, Takeda Y. C18 Acetylenic Fatty Acids of *Ximenia americana* with potential pesticidal activity. *J Agr Food Chem* 2000; 48: 1872-1874. <https://doi.org/10.1021/jf990550k>
13. Visuthikosol V, Chowchuen B, Sukwanarat Y, Sriurairatana S, Boonpucknavig V. Effect of *Aloe vera* gel to healing of burn wound a clinical and historic study. *J Med Assoc Thai* 1995; 78: 403-9. PMID:7561562