Available online on 15.07.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal

Open access to Pharmaceutical research
 This is an open access article distributed under the terms of the Creative Commons Attribution-Non
 Commercial Share Alike 4.0 License which permits unrestricted non commercial use,
 provided the original work is properly cited

Volume 5, Issue 3, 2020

Open Access

RESEARCH ARTICLE

BACTERIAL CAUSES AND ANTIMICROBIAL SENSITIVITY PATTERN OF EXTERNAL OCULAR INFECTIONS IN SELECTED OPHTHALMOLOGY CLINICS IN SANA'A CITY

Essam Yahya A Alshamahi¹, Huda Zaid Al-Shami², Hassan A. Al-Shamahy²,
 Yaser Ali Musawa²

¹Ophthalmology Department, Faculty of Medicine and Health Sciences, Sana'a University, Republic of Yemen.

²Medical Microbiology and Clinical Immunology Department, Faculty of Medicine and Health Sciences, Sana'a University, Republic of Yemen.

ABSTRACT

Objectives: The aim of the current study was to reveal the bacterial profile and pattern of sensitivity to antibiotics for external ocular infections for patients who attended selected ophthalmology clinics in the city of Sana'a.

Methods: A cross-sectional study design was used from September 2016 to October 2017 where a total of 197 patients with infection of external eye were included in the study which included conjunctivitis, keratitis, blepharitis and Blepharoconjunctivitis. Samples were collected and transferred to the National Center of Public Laboratories (NCPHL), in Sana'a. Possible bacterial pathogens have been isolated and identified using regular laboratory techniques, and microbial sensitivity testing has been carried out using a disc diffusion method.

Results: A total of 197 ocular samples were obtained for microbiological evaluation, of these 146 (74.1%) have bacterial growth. Bacteria of Gram positive accounted for 52.1% and the prevalent isolation was *S. aureus* (30.1%). Gram negative bacteria made up 47.9% and the predominant isolation was *Pseudomonas aeruginosa* (26.7%). The majority of Gram-positive bacteria were sensitive to ciprofloxacin (90% - 100%), vancomycin (86% - 100%) and Gram-negative isolates sensitive for amikacin (100%) and ciprofloxacin (63% - 100%).

Conclusion: These results revealed that Gram-positive bacteria were the generally common bacteria isolated from infections of external eye and were more susceptible to vancomycin and ciprofloxacin while Gram-negative isolates were more susceptible to ciprofloxacin and amikacin. The high rate of resistance for most antibiotics in Yemen, leaves ophthalmologists with very few options of drugs to treat eye infections. Large-scale ongoing studies in the future should also be conducted in order to monitor the antimicrobial resistance of the external ocular bacterial isolates.

Keywords: antimicrobial sensitivity, bacterial causes, external ocular infections, Sana'a, Yemen.

Article Info: Received 2 May 2020; Revised 6 June; Accepted 4 July, Available online 15 July 2020

Cite this article-

Alshamahi EYA, Al-Shami HU, Al-Shamahy HA, Musawa YA. Bacterial causes and antimicrobial sensitivity pattern of external ocular infections in selected ophthalmology clinics in Sana'a city. Universal Journal of Pharmaceutical Research 2020; 5(3):12-16.

DOI: <https://doi.org/10.22270/ujpr.v5i3.409>

Address for Correspondence:

Prof. Hassan A. Al-Shamahy, Medical Microbiology and Clinical Immunology Department, Faculty of Medicine and Health Sciences, Sana'a University, Republic of Yemen, Tel: +967-1-239551, E-mail: shmahe@yemen.net.ye

INTRODUCTION

Pathogenic microorganisms cause ocular infections as a result of virulence and low host resistance in some circumstances such as living conditions, personal hygiene, socioeconomic status, low immunity status, and other related factors^{1,2}. The conjunctiva, the lid and the cornea are the frequently affected areas of the eye^{1,2}. Bacteria are one of the main causative agents that cause eye infections, which may lead to blindness. Thus, an immediate treatment of a serious bacterial eye infection that threatens the cornea is needed². For

precise antibacterial treatment, isolation and identification of bacterial pathogens along with an antibiotic sensitivity spectrum is required³. Because there is a global problem with the appearance of bacterial resistance to topical antimicrobial agents that are effected by pathogen properties and antibiotic prescribing practices including systemic antibiotic use and general health care guidelines^{4,5}. This growing resistance increases the risk of treatment failure with potentially severe consequences^{6,7}. Bacterial etiology and sensitivity, as well as patterns of resistance, may

vary according to geographical and regional location⁶⁻⁸. Hence, recent information is vital for ophthalmologists for proper antimicrobial therapy^{1,4,6,7}. In Yemen, there was no previous study conducted on external ocular infections and patterns of sensitivity to antibiotics before this study. The study carried out by Al-Shamhi and others studied epidemiology and the diagnosis of corneal ulcers in the city of Sana'a, could be considered as part of this study as it only focused on corneal infection⁹. Due to the high rate of drug resistance to antibiotics in medicine in Yemen¹⁰, the ophthalmologists is left with a very few options of drugs for the treatment of ocular infections. Hence, knowledge of the causative agents of this infection is essential to proper case management. Bacterial sensitivity to many antimicrobial agents changes from location to location and in the same place from time to time^{6,8}. Consequently, the changing spectrum of microorganisms concerned in eye infections and the emergence of acquired microbial resistance determine the need for continuous monitoring to guide experimental treatment^{6,7,11}. The experimental choice of effective treatment has become more complicated as ocular pathogens are becoming increasingly resistant to commonly used antibiotics⁷. Regarding the study area in Yemen, there is a scarcity or lack of published data on the spectrum of etiological agents responsible for external eye infections. Thus, the purpose of this study was to determine the spectrum of bacterial etiology for external ocular infections, and to assess the susceptibility of these bacterial ocular isolates to in vitro antibiotics regularly prescribed among patients with external eye infections in Yemen.

MATERIALS AND METHODS

This cross-sectional study included 197 eye specimens for microbiological assessment of patients clinically diagnosed with external eye infections such as keratitis, conjunctivitis, blepharitis and blepharo-conjunctivitis in selected ophthalmology clinics in Sana'a between September 2016 to October 2017. Patients diagnosed clinically with external eye infections, with informed consent taking, were included in the study. Excluding patients with viral conjunctivitis, trachoma, viral keratitis, peripheral ulcerative keratitis, allergic and severe eye trauma, current eye surgery, and patients

who received antimicrobial therapy within two weeks before the requirement. All patients were examined on a slit lamp biomicroscope and the infectious diseases included in this study were clinically diagnosed by a group of ophthalmologists. After detailed eye examination using standard techniques¹², samples from the eyelid, conjunctiva, and cornea were collected by ophthalmologists. Immediately obtained eye samples were inoculated in blood agar, chocolate agar, selective media for MNYC [if the newborn patient and *N. gonorrhea* were suspected], as well as the Loeffler serum slope of the *Moraxella* infection (all culture media from Difco Laboratories USA). Then the plates and tubes were incubated in appropriate conditions. Possible bacterial pathogens were isolated and identified using standard laboratory techniques, and microbial sensitivity testing was performed by a disc diffusion method¹³. The following antimicrobials were used with their respective concentration (Difco Laboratories, USA): Ceftriaxone (CRO, 30 µg), amikacin (AK, 30 µg), gentamicin (CN, 10 µg), ciprofloxacin (CIP, 5 µg), penicillin (P, 10U), tetracycline (TE, 30 µg), erythromycin (E, 15 µg), doxycycline (DO, 30 µg), chloramphenicol (C, 30 µg), trimethoprim-sulphamethoxazole (SXT, 1.25/23.75 µg), and vancomycin (VA, 30 µg).

Table 1: The age and gender distribution of patients with external ocular infection in selected hospitals and eye clinics in Sana'a city, Yemen.

Characters	Male (n= 197)	
	No.	%
Sex		
Male	121	61.4
female	76	38.6
Age group		
≤ 15 years	47	23.9
16 – 25 years	31	15.7
26 – 35 years	42	21.3
36 – 45 years	29	14.7
≥ 46 years	48	24.4
Total	197	
Mean age	29.4 years	
S D	9.5 years	
Min	1 month	
Max	80 years	

Table 2: Distribution of bacterial isolates of external ocular infection in selected hospitals and eye clinics in Sana'a city, Yemen.

Bacterial isolates	Positive for bacterial growth N=146	
	No	%
Gram positive bacteria	76	52.1
<i>Staphylococcus aureus</i>	44	30.1
<i>Beta-hemolytic streptococcus</i>	9	6.2
<i>Streptococcus pneumonia</i>	11	7.5
<i>Coagulase negative staphylococci</i>	12	8.2
Gram negative bacteria	70	47.9
<i>Haemophilus influenzae</i>	13	8.9
<i>Pseudomonas aeruginosa</i>	39	26.7
<i>Moraxella lacunata</i>	5	3.4
<i>Escherichia coli</i>	11	7.5
<i>Proteus species</i>	2	1.4
Total n=197	146	74.1

Table 3: Antibiotic susceptibility test of Gram-positive isolates from external ocular infections in selected hospitals and eye clinics in Sana'a city, Yemen.

Antibiotic	<i>S. aureus</i> N=44		<i>S. pneumoniae</i> N=11		CoNs N=12		<i>Beta hemolytic streptococci</i> N=9		Total N=76	
	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)
Amikacin (30 µg)	42(95.5)	2(4.5)	2(18.2)	9(81.8)	10(83.3)	2(16.7)	3(33.3)	6(66.7)	57(75)	19(25)
Penicillin (30 µg)	1(2.3)	43(97.7)	9(81.8)	2(18.2)	5(41.7)	7(58.3)	6(66.7)	3(33.3)	21(27.6)	55(72.4)
Vancomycin (30 µg)	38(86.4)	6(23.6)	11(100)	0(0.0)	12(100)	0(0.0)	9(100)	0(0.0)	70(92.1)	6(7.9)
Erythromycin (15 µg)	26(59)	18(41)	9(81.8)	2(18.2)	8(66.7)	4(33.3)	7(77.8)	2(22.2)	50(65.9)	26(34.1)
Trimethoprim-sulphamethoxazole (1.25/23.75 µg)	32(72.7)	12(27.3)	5(45.5)	6(54.5)	8(66.7)	4(33.3)	7(77.8)	2(22.2)	52(68.4)	24(31.6)
Chloramphenicol (30 µg)	23(52.3)	21(47.7)	11(100)	0(0.0)	7(58.3)	5(41.7)	8(88.9)	1(11.1)	49(64.5)	27(35.5)
Gentamicin (10 µg)	40(90.9)	4(9.1)	3(27.3)	8(72.7)	6(50)	6(50)	5(55.6)	4(44.4)	54(71)	22(29)
Tetracycline (30 µg)	30(68.2)	14(31.8)	3(27.3)	8(72.7)	9(75)	3(25)	3(33.3)	6(66.7)	45(59.2)	31(40.8)
Doxycycline (30 µg)	34(77.3)	10(22.7)	10(90.9)	1(9.1)	11(91.7)	1(8.3)	8(88.9)	1(11.1)	63(83)	13(17)
Ceftriaxone (30 µg)	30(68.2)	14(31.8)	9(81.8)	2(18.2)	9(75)	3(25)	8(88.9)	1(11.1)	56(73.7)	20(26.3)
Ciprofloxacin (5 µg)	42(95.5)	2(4.5)	10(90.9)	1(9.1)	12(100)	0(0.0)	9(100)	0(0.0)	73(96)	3(4)

RESULTS

A total of 197 external ocular infection patients (121 - 61.4%, male and 76 - 38.6%, female) were enrolled in this study. The most frequent age groups were ≤ 15 years (23.9%), and age group ≥ 46 years (24.4%); while young adult groups were less frequent (Table 1). Bacterial growth yielded on 146 (74.1%) while 51 (25.9%) were negative for bacterial culture (Table 2). The isolates in 146 patients with external ocular infections were Gram positive bacteria (52.1 %), the predominant species of Gram positive was *S. aureus* (30.1%), while *beta hemolytic streptococci* counted 6.2%, *S. pneumoniae* was 6.2%, and CoNs was 8.2%. Gram negative isolates counted for 47.9%, the

predominant Gram negative bacteria was *P. aeruginosa* (26.7%), while other species were less frequent e.g. *E. Coli* (7.5%), *Moraxella* species (3.4%), *H. influenzae* (8.9%) and *Proteus* species was 1.4% (Table 2). Table 3 shows the sensitivity of Gram-positive bacteria. Most of Gram-positives showed resistance against penicillin up to 97%; but they were highly susceptible to ciprofloxacin (96%), vancomycin (92%), doxycycline (83%), tetracycline (59.2%), ceftriaxone (73.7%), erythromycin (92%). Table 4 illustrates the susceptibility of Gram-negative bacteria. The Gram-negative bacteria and showed high rate of susceptibility to amikacin (100%) and gentamicin (89.7%).

Table 4: Antibiotic susceptibility test of Gram-negative isolates from external ocular infections in selected hospitals and eye clinics in Sana'a city, Yemen.

Antibiotics	<i>H. influenzae</i> N=13		<i>Pseudomonas aeruginosa</i> N=39		<i>E.coli</i> N=11		<i>Moraxella lacunata</i> , N=5		Total N=68	
	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)	S No (%)	R No (%)
Amikacin (30 µg)	13(100)	0(0.0)	39(100)	0(0.0)	11(100)	0(0.0)	5(100)	0(0.0)	68(100)	0(0.0)
Erythromycin (15 µg)	9(69.2)	4(30.8)	NT	NT	NT	NT	NT	NT	9(69.2)	4(30.8)
Trimethoprim-sulphamethoxazole (1.25/23.75 µg)	2(15.4)	11(84.6)	8(20.5)	31(79.5)	2(18.2)	9(81.8)	1(20)	4(80)	13(19.1)	55(80.9)
Chloramphenicol (30 µg)	9(69.2)	4(30.8)	23(59)	16(41)	4(36.4)	7(63.6)	5(100)	0(0.0)	41(60.3)	27(39.7)
Gentamicin (10 µg)	13(100)	0(0.0)	38(97.4)	1(2.6)	6(54.5)	5(45.5)	4(80)	1(20)	61(89.7)	7(10.3)
Tetracycline (30 µg)	10(76.9)	3(23.1)	10(25.6)	29(74.4)	8(72.7)	3(27.3)	4(80)	1(20)	32(47)	36(53)
Doxycycline (30 µg)	11(84.6)	2(15.4)	21(53.8)	18(46.2)	7(63.6)	4(36.4)	4(80)	1(20)	43(63.2)	25(36.8)
Ceftriaxone (30 µg)	12(92.3)	1(7.7)	30(76.9)	9(23.1)	5(45.5)	6(54.5)	4(80)	1(20)	51(75)	17(25)
Ciprofloxacin (5 µg)	11(84.6)	2(15.4)	35(89.8)	4(10.2)	7(63.6)	4(36.4)	5(100)	0(0.0)	58(85.3)	10(14.7)

NT= Not tested, S=sensitive, R=resistance

DISCUSSION

In wide-ranging, the eye isolates recognized in the current study were comparable to those of many other studies performed in different regions. Although the major bacteria recognized to cause external eye infections around the world is *S. aureus*^{1,2,8,14}. The most common isolates in the current study was *S. aureus* (30.1%) followed by *P. aeruginosa* (26.7%). Similar studies performed in India^{1,8}, Nigeria^{2,15}, Gondar¹⁶, and

Ethiopia^{14,17} also indicated that *S. aureus* is a predominant eye isolate. On the other hand, some other studies have reported that *S. aureus* is the first but has reported *E. coli*^{9,15}, *S. albus*², *S. pneumoniae*¹⁸ as the second common bacterial isolation not *P. aeruginosa* such as the current study. The predominance of *P. aeruginosa* in our study, which differs from previous studies, can be supported by finding similar studies conducted in Sudan¹⁸, Australia¹⁹, Malaysia²⁰, India²¹ and Thailand²². These results can be explained by the

fact that as part of the eye's natural flora, *Pseudomonas* grow better in the eye than any recognized culture media and cause infection when mechanical shock to the corneal epithelium occurs, also, it produces external toxins A, which cause tissue necrosis leading to corneal ulceration^{2,12}. The present study showed fewer isolates of intestinal bacteria (*E. coli* =7.5%; *Proteus* =1.4%) when compared to a similar study performed in Nigeria¹⁵, and Gondar¹⁶ where variable *Enterobacteriaceae* were more common isolates from external eye infections. This low number of intestinal bacteria in our study may be due to decreased in hand-faecal contamination and/or increased access to safe drinking water sources in the study area². Laboratory-based resistance and sensitivity may not reflect the true clinical resistance and response to the antibiotic due to drug penetration and host factors⁸. On the other hand, these findings afford data that allows the doctor to make a rationale-based decision in choosing a primary regimen for ocular pathogens¹. Based on the results of the sensitivity test in the current study, most Gram-positive bacteria were susceptible to ciprofloxacin (96%) followed by vancomycin (92.1%). Vancomycin coverage against *S. aureus* and CoNS was 86.4% and 100%, respectively. This result corresponds to the study carried out in India⁸. In contrast, a study in Iran¹¹ reported low coverage of vancomycin against *S. aureus*. Gentamicin covered against 71% of Gram-positive isolates and obtained high coverage against *S. aureus* (90.9%) This finding is similar with studies conducted in Nigeria^{2,14}, Iran¹¹, and India^{8,21}. Though, this study showed low gentamicin coverage against CoNS (50%) compared to a study carried out in Nigeria^{2,14}, and India²¹. Gentamicin coverage for *Pseudomonas aeruginosa* was 97.4%. *P. aeruginosa*, which makes up 26.7% of all isolated bacteria, was highly sensitive to amikacin (100%), ciprofloxacin (89.8%), ceftriaxone (76.9%), doxycycline (53.8%), and chloramphenicol (59%). These results were reported for ciprofloxacin from studies conducted in Saudi Arabia²⁴ and Nigeria^{14,25}. On the other hand, the study in India²¹ described low coverage of ciprofloxacin for *P. aeruginosa*. The gentamicin coverage against Gram-negative bacteria in this study was 89.7%. This can be compared to similar studies conducted in Nigeria² and Iran¹¹. In spite of this, the study carried out in India²⁶ indicated that gentamicin coverage for Gram-negative bacteria including *P. aeruginosa* was low. Tetracycline coverage against Gram-positive bacteria was 59.2%. This result is comparable with the study carried out in Iran and Nigeria^{11,14}. Tetracycline coverage against Gram-positive bacteria was 59.2%. This result is comparable with the study carried out in Iran and Nigeria^{11,14}. The majority of Gram-negative bacteria (72%) appeared to be resistant to penicillin. In spite of this, coverage of penicillin against *S. pneumoniae* was high in the current (81.8%). This is comparable to studies done in Iran¹¹ and Nigeria². Amikacin has high coverage against *S. aureus* (95.51%) and CoNS (83.3%). This is regular with studies done in Iran¹¹ and India^{8,21}. There is an increase in the resistance of studied antibiotics against isolated bacteria in the current study as in other

studies elsewhere, the emergence of bacterial resistance due to pathogen properties and antibiotic prescribing preparations including the extensive use of systemic antibiotics and health care guidelines^{5,6,27}. Other causal factors may comprise an improper dose regimen, mistreatment of antibiotics for viral infections and other non-bacterial infections, and a long period of treatment rather than in the least globalization and migration⁶. In Yemen, it is a widespread practice that antibiotics can be acquired without a prescription, leading to misuse of antibiotics. This may contribute to the emergence and spread of antimicrobial resistance^{16,17}. Other factors may include substandard quality or substandard antimicrobial drugs, increased use of a specific antimicrobial agent, contaminated food, poor sanitation, and fecal contamination from animals or humans^{2,5,17}. As a result of patterns of bacterial sensitivity to many antimicrobial agents, they may vary from place to place and in the same place from time to time^{7,8,23}.

CONCLUSION

These results revealed that Gram-positive bacteria were the most common bacteria isolated from external eye infections and were more susceptible to vancomycin and ciprofloxacin while Gram-negative isolates were more susceptible to ciprofloxacin and amikacin. The high rate of resistance for most antibiotics in Yemen, leaves ophthalmologists with very few options of drugs to treat eye infections. Mono and gatifloxacin, fusidic acid, tobramycin, neomycin that are used as eye drops were not included in the tested antibiotics because they were not available in our laboratory during the study. We usually only test antibiotics used for systemic infections. Extensive future studies should also be conducted in order to monitor antimicrobial resistance including topical antibiotics such as mono, and gatifloxacin, fusidic acid, tobramycin, and neomycin. On the other hand external fungal eye infections should also be studied.

ACKNOWLEDGEMENTS

The authors would like to acknowledge Ministry of Health and Population, Sana'a, Yemen and the National Center of Public Health Laboratories (NCPHL), Ministry of Health and Population, Sana'a, Yemen for their support and provided working space and materials.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

AUTHOR'S CONTRIBUTION

This research work is part of the National Center for Public Health Laboratories (NCPHL) and MSc project. The first author did clinical work and the second and third authors did the lab. The corresponding author (HAA) supervised laboratory works, reviewing and editing the research.

REFERENCES

1. Ramesh S, Ramakrishnan R, Bharathi MJ, Amuthan M, Viswanathan S. Prevalence of bacterial pathogens causing

- ocular infections in South India. *Indian J Pathol Microbiol* 2010; 53(2):281-286.
<https://doi.org/10.4103/0377-4929.64336>
2. Ubani UA. Bacteriology of external Ocular Infections in Aba, South Eastern Nigeria. *Clin Exp Optom* 2009; 92(6):482-9.
<https://doi.org/10.1111/j.1444-0938.2009.00425.x>
 3. Chung JL, Seo KY, Yong DE, Mah FS, Kim T, Kim EK, *et al.* Antibiotic Susceptibility of Conjunctival Bacterial Isolates from Refractive Surgery Patients. *Ophthalmol* 2009; 116(6):1067- 74.
<https://doi.org/10.1016/j.ophtha.2008.12.064>
 4. Lee K, Lee H, Kim M. Two cases of corneal ulcer due to methicillin-resistant *Staphylococcus aureus* in high risk groups. *Korean J Ophthalmol* 2009; 24(4):240-4.
<https://doi.org/10.3341/kjo.2010.24.4.240>
 5. Joseph S, Bertino JR. Impact of antibiotic resistance in the management of ocular infections: the role of current and future antibiotics. *Clin Ophthalmol* 2009; 3:507-21.
<https://doi.org/10.2147/ophth.s5778>
 6. Sharma S. Antibiotic resistance in ocular bacterial pathogens. *Indian J Med Microbiol* 2011; 29:218-222.
<https://doi.org/10.4103/0255-0857.83903>
 7. Brown L. Resistance to ocular antibiotics: an overview. *Clin Exp Optom* 2007; 90(4):258-62.
<https://doi.org/10.1111/j.1444-0938.2007.00154.x>
 8. Bharathi MJ, Ramakrishnan R, Shivakumar C, Meenakshi, R, Lionaraj D. Etiology and antibacterial susceptibility pattern of community-acquired bacterial ocular infections in a tertiary eye care hospital in south India. *Indian J Ophthalmol* 2010; 58(6):497-507.
<https://doi.org/10.4103/0301-4738.71678>
 9. Alshamahi EYA, Al Shamahy HA Al Nahary AA, AL Magrami RTF, *et al.* Epidemiology and Aetiological Diagnosis of Corneal Ulceration in Sana'a City, Yemen. *W J Ophthalmol Vision Res* 2019; 2(5):1-6.
<https://doi.org/10.33552/WJOVR.2019.02.000550>
 10. Al-Safani AMA, Al-Shamahy HA, Al-Moyed KA. Prevalence, antimicrobial susceptibility pattern and risk factors of MRSA isolated from clinical specimens among military patients at 48 medical compound in Sana'a city Yemen. *Universal J Pharm Res* 2018; 3(3): 40-44.
<https://doi.org/10.22270/ujpr.v3i3.165>
 11. Khosravi AD, Mehdinejad M, Heidari M. Bacteriologic Findings in Patients with Ocular Infection and Antibiotic Susceptibility Patterns of Isolated Pathogens. *Singapore Med J* 2007; 48(8):741-3. *PMID: 17657382*
 12. Byrne KA, Burd E, Tabbara K, Hyndiuk R. *Diagnostic Microbiology and Cytology of the Eye*. Butterworth-Heinemann, Boston 1995; 40-2. *PMCID: PMC500739*
 13. Cheesbrough M. *District laboratory practice in tropical countries*. Cambridge, New York 2006; 132-5.
 14. Tesfaye T, Beyene G, Gelaw Y, *et al.* Bacterial profile and antimicrobial susceptibility pattern of external ocular infections in Jimma University Specialized Hospital, Southwest Ethiopia. *American J Infect Dis Microbiol* 2013; (1):13-20. <https://doi.org/10.12691/ajidm-1-1-3>
 15. Esenwah E. Isolation and identification of the microorganisms most prevalent in external eye infections as seen in an eye clinic in Owerri. *J Nig Opto Assoc* 2005; 12:6-9. <https://doi.org/10.4314/jnoa.v12i1.64449>
 16. Anagaw B, Biadlegne F, Belyhun, Y, Mulu A. Bacteriology of ocular infections and antibiotic susceptibility pattern in Gondar University Hospital, North west Ethiopia. *Ethiop Med J* 2011; 49(2):117-23. *PMID: 21796911*
 17. Alemayehu N. Pattern of Microbial Agents of External Ocular Infections in Federal Police Hospital and Minilik II Memorial Hospital. AAU: MSc thesis. 2004.
 18. Bataineh H, Hammory Q, Khatatba A. Bacterial Keratitis: Risk Factors and Causative Agents. *Sudan J Med Sci* 2008; 3(1):6-10. <https://doi.org/10.4314/sjms.v3i1.38503>
 19. Green M, Apel A, Stapleton F. Risk factors and causative organisms in microbial keratitis. *Cornea* 2008; 27(1):22-27. <https://doi.org/10.1097/ICO.0b013e318156caf2>
 20. Norina TJ, Raihan S, Bakiah S, Ezanee M, Liza-Sharmini AT, Hazzabah WH. Microbial keratitis: aetiological diagnosis and clinical features in patients admitted to Hospital Universiti Sains Malaysia. *Singapore Med J* 2008; 49(1):67-71. *PMID: 18204773*
 21. Biradar S, Chandrashekhara DK, Gangane R, *et al.* Spectrum of microbial keratitis and antimicrobial susceptibility at tertiary care teaching hospital in north Karnataka. *Int J Pharm Biomed Res* 2012; 3(2):117-20.
 22. Tananuvat N, Sienglew S, Ausayakhun S. Microbial keratitis leading to admission at Maharaj Nakorn Chiang Mai hospital. *Chiang Mai Med Bull* 2004; 43(3):93-103.
 23. Tobbara KF, Hyndiuk RA. *Infections of the Eye*, Little, Brown and Company, New York, 1995, 55-7.
 24. Al-Zahrani SHM. Bacteria isolated from contact and non contact lens and antibiotic susceptibility patterns of isolated *Pseudomonas aeruginosa*. *Afr J Microbiol Res* 2012; 6(47):7350-6. <https://doi.org/10.5897/AJMR12.1134>
 25. Ogbolu DO, Alli OT, Ephraim IE, Olabiyi FA, Daini OA. *In-vitro* efficacy of antimicrobial agents used in the treatment of bacterial eye infections in Ibadan, Nigeria. *Afr J Clin Exper Microbiol* 2011; 12:124-127.
<https://doi.org/10.4314/ajcem.v12i3.7>
 26. Akpek EK, Gottsch JD. Immune defense at the ocular surface. *Eye* 2003; 17(8):949-56.
<https://doi.org/10.1038/sj.eye.6700617>
 27. Cavuoto K, Zutshi D, Karp CL, Miller D, Feuer W. Update on Bacterial Conjunctivitis in South Florida. *Ophthalmol* 2008; 115(1):51-6.
<https://doi.org/10.1016/j.ophtha.2007.03.076>