Available online on 15.07.2020 at <http://ujpr.org>
Universal Journal of Pharmaceutical Research
 An International Peer Reviewed Journal
 Open access to Pharmaceutical research

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial Share Alike 4.0 License which permits unrestricted non commercial use, provided the original work is properly cited

Volume 5, Issue 3, 2020

Open Access

REVIEW ARTICLE

INFLAMMATORY BOWEL DISEASE: CLINICAL FEATURES AND MANIFESTATIONS BEYOND THE BOWEL

Maged Almezgagi^{1,2,7} , Yu Zhang² , Fadhl Al-shaebi³ , Mueataz A. Mahyoub⁴ , ALariqi Reem⁵ , Muhammad Shoaib¹ , Emad Shamsan² , Mohammed Gamah^{1,2} , Kamal Hezam⁶ , Abdul Baset Abbas⁷ , Wang Xiaozhou¹ , Shen Xiangqun¹ , Ma Qianqian¹ , Ying Han¹ , Ruhan Jia¹ , Wei Zhang^{1,2}

¹The Key Laboratory of high-altitude medical application of Qinghai Province, Qinghai Xining 810001, China.

²Department of Immunology, of Qinghai University, Qinghai Xining 810001, China.

³Department of Immunology, Hebei Medical University, Shijiazhuang, Hebei 050017, China.

⁴Department of Gastroenterology, Second Affiliated Hospital, Medical College of Xi'an Jiaotong University, Xi'an 710004

⁵Department of Microbiology, Harbin Medical University, Harbin 150081, China.

⁶Nankai University School of Medicine, Tianjin 300071, PR China.

⁷Department of Medical Microbiology, Faculty of Sciences, Ibb University, Ibb City 70270, Yemen.

ABSTRACT

Inflammatory bowel disease (IBD) encompasses a spectrum of diseases, with Crohn's disease (CD) and ulcerative colitis (UC) representing the two broadest subtypes of IBD. Multiple extraintestinal manifestations (EIMs) are more frequent in (IBD); 5% – 50% of the patients might be affected. The most often implicated sites of manifestations are musculoskeletal and dermatological structures. However, while some symptoms like peripheral arthritis and erythema nodosum correlate with IBD progression, others have their own course of disease like axial arthropathy, gangrenosis of the pioderma and primary sclerosing cholangitis. This review would provide a summary of the most frequent EIMs and their prevalence.

Keywords: Crohn's disease, inflammatory bowel disease, Ulcerative colitis.

Article Info: Received 22 April 2020; Revised 1 June; Accepted 4 July, Available online 15 July 2020

Cite this article-

Almezgagi M, Zhang Yu, Al-shaebi F, Mahyoub MA, Reem A, Shoaib M, Shamsan E, Gamah M, Hezam K, Abbas AB, Xiaozhou W, Xiangqun S, Qianqian Ma, Han Y, Jia R, Zhang W. Inflammatory bowel disease: clinical features and manifestations beyond the bowel. Universal Journal of Pharmaceutical Research 2020; 5(3):68-74.

DOI: <https://doi.org/10.22270/ujpr.v5i3.422>

Address for Correspondence:

Wei Zhang, The Key Laboratory of high-altitude medical application of Qinghai Province, Qinghai Xining 810001. Department of Immunology, of Qinghai University, Qinghai Xining 810001, China. E-mail: zw0228@qhu.edu.cn

INTRODUCTION

IBD is a chronic disease which relapses immune-mediated. The two main IBD subtypes are UC and CD¹. UC is an idiopathic, chronic condition in the colon, most frequently affecting adults between 30 and 40 years of age with disabilities^{2,3}. It is characterized by relapse and remittance of inflammation in mucosal tissues, beginning in the rectum then spreading to proximal colon segments. UC marked by cycles of recovery and cycles of recurrent, the latter, frequently presenting with the abdominal pain, vomiting, rectal bleeding, weight loss and malaise, is responsible for the most common of the disease problem^{4,5}. The risk for UC newly diagnosed patients is between 10% and 35% for five years and ultimately the long-time risk of colorectal cancer is enhanced by extensive and

persistent inflammatory condition⁶. The control of symptoms such as increasing in the bowel movement and rectal hemorrhage was a controlled problem⁵. The introduction of standardized clinical scores, such as the Truelove and Witts criteria⁷ and the Mayo score⁵, allowed for a more accurate evaluation of the disease and, while they are sometimes used in clinical trials^{8,9}, have not yet been validated. This strategy, aimed at regulating and alleviating the effects of inflammation, did not target the inflammatory activity itself. In comparison, there is a major correlation between clinical IBD and other disorders, such as irritable bowel syndrome (IBS) or infectious diarrhea¹⁰, and some authors recorded a long-term recovery of UC patients with IBS-like symptoms (abdominal pain, elevated stool frequency) two to three times more

frequency than controls¹⁴. In contrast, others observed high stool frequency in up to 27% of patients with full endoscopic and histological healing, indicating a potential explanation for non-inflammatory functional intestinal damage¹⁵. Finally, in a systematic study of clinical trials, clinical recovery during the placebo reached up to 15%¹⁶. However, there is growing evidence that clinical improvement without mucosal healing (MH) is not associated with reduced hospitalization or colectomy rates over the years^{17,18}. Inflammatory markers such as erythrocyte sedimentation rate, fecal marker calprotectin and serum markers C-reactive protein are other desirable choices for tracking UC patients¹¹. Further studies are required to explain adequate surveillance strategies and cut-off rates before it is widely used in clinical practice. Mucosal inflammation is a central feature of both CD and UC but, unlike CD, a disease with both strict and penetrating phenotypes, the development of the disease is limited to the UC mucosa¹². Therefore, it is no surprise that MH will prove an appealing target when addressing UC patients regardless of the severity of the disorder, inflammatory biomarkers or clinical

presentation. Extensive research has been published over the past decade supporting the value of histological healing^{13,14} as it has shown great correlation with decreased risk of relapse¹⁵ and hospitalization¹⁶. In some studies, histological healing may be involved in the description of MH beside the findings of endoscopy¹⁶. The therapy is aimed at inducing and sustaining clinical and endoscopic remission. Current treatment choice for UC includes aminosaliclates, such as corticosteroids (including systemic corticosteroids such as hydrocortisone or prednisolone, and topical corticosteroids such as budesonide), mesalamine (5-aminosalicylic acid; 5-ASA) in both rectal and oral preparations, sulfasalazine, thiopurines (6-mercaptopurine and azathioprine), calcineurin inhibitors (tacrolimus and cyclosporine), anti-tumor necrosis factor (TNF)-a drugs (including golimumab, adalimumab and infliximab), methotrexate and more newly, the anti-integrin drug vedolizumab¹⁶. This review summarizes the most common EIMs and its clinical features and their prevalence and suggested management.

Table 1: Extraintestinal manifestations of IBD.

Gastrointestinal	Primary sclerosing cholangitis (PSC), PSC–autoimmune “hepatitis overlap syndrome”, Drug-induced hepatitis, Hepatic steatosis, Hepatic abscess, Portal vein thrombosis Pancreatitis, Immunoglobulin G4 (IgG4)–associated cholangitis, Primary biliary cirrhosis, Cholelithiasis, Autoimmune pancreatitis
Urinary	Enterourinary fistulas, Obstructive uropathy, Nephrolithiasis
Musculoskeletal	Arthritis: isolated joint involvement, ankylosing spondylitis, Hypertrophic osteoarthropathy: periostitis, clubbing Other: aseptic polymyositis, necrosis
Pulmonary	Large airways disease, Pneumonia
Cardiac	Congestive heart failure
Ocular system	Episcleritis, uveitis/iritis, scleromalacia, retinalvascular disease, corneal ulcers, Conjunctivitis, Orbital pseudotumor
Dermatologic/Oral	Reactive lesions: pyoderma gangrenosum, erythema nodosum, necrotizing vasculitis, aphthous ulcers; Specific lesions: fistulas, fissures, drug rashes, oral Crohn disease; Nutritional deficiencies: purpura, acrodermatitis, enteropathica, hair loss, glossitis, brittle nails; Associated diseases: amyloidosis, psoriasis, vitiligo
Hematologic	Anemia, hyperhomocysteinemia
Metabolic system	Delayed sexual maturation, growth retardation in children and adolescents, osteoporosis/osteopenia

2. Extraintestinal manifestations

CD and UC also lead to extraintestinal manifestations (Table. 1), as shown in 25 to 40 % of IBD patients⁴⁵. Almost any part in the body at risk, but the primary manifestations are symptoms affecting the joints, liver, skin and eyes. Existing one extraintestinal symptoms raises the likelihood that another will develop⁴⁶. The treatment of underlying gut inflammation leads to the symptoms of concurrent disease activity such as erythema nodosum, episcleritis peripheral and arthritis. IBD can be associated with a specific disease, for example, IBD is correlated with primary sclerosing cholangitis (PSC); 5% to 10% have CD and 75% of PSC patients have UC. Nevertheless, only 2% of CD patients develop PSC and about 5% of UC patients,

respectively⁴⁷. IBD patients who have persistent diarrhea, usually with mucous and blood. For UC, symptom period may vary and appear to be more indolent, lasting weeks to months¹⁷⁻¹⁹. Extraintestinal manifestations in 21%-47% of patients with IBD are reported¹. Most extraintestinal manifestations are not well known for pathogenesis. In recent years, however, significant advances in the genetic basis of IBD have occurred with the advent of genomewide interaction studies. In particular, 99 susceptibility loci or genes have been reported to date (47 in ulcerative colitis and 71 in CD)²⁰. Many organs may be involved with extraintestinal manifestations in IBD patients. Extraintestinal symptoms in joints, skin, hair have been correlated with the level of bowel inflammation, but

cardiothoracic and gastrointestinal (hepatobiliary) desorption generally did not have that correlation¹.

2.1 IBD Manifestations of gastrointestinal

Pancreatic and hepatobiliary damage consider the most severe extraintestinal manifestations in IBD patients. While the relations between many liver disorders and IBD are well known, other associations are far more²¹.

2.1.1 Primary Sclerosing Cholangitis

In the IBD setting, PSC is the most common hepatobiliary manifestation²⁰. PSC symptoms are progressive inflammation in the biliary tree, obliterative fibrosis and death, resulting in biliary

fibrosis, cirrhosis and probable hepatic failure²². The PSC-IBD relationship was first identified by Smith and Loe in 1965²³.

2.1.2 PSC–Autoimmune (Hepatitis Overlap Syndrome)

In IBD patients, especially in ulcerative colitis patients, an association between autoimmune hepatitis and PSC has been reported. Several case reports of IBD patients who were initially diagnosed with autoimmune hepatitis later established PSC histological evidence²⁴.

Figure 1: Pancreatic manifestations accompanied by IBD.

2.1.3 Cholangiocarcinoma

The emergence of cholangiocarcinoma, a crippling malignancy with an exceedingly poor prognosis, is a feared complication of PSC. Though cholangiocarcinoma does not result directly from IBD, the literature has well established a significant relation between cholangiocarcinoma and PSC¹. PSC patients tend to show cholangiocarcinoma earlier than intermittent cholangiocarcinoma²⁵. With a lifetime incidence of 5%-15%, PSC patients have a substantially higher chance of cholangiocarcinoma progress²⁶.

2.1.4 Drug-induced Hepatitis

Although hepatobiliary disorders described above share common IBD pathogenesis; many medicines used to treat IBD can cause liver toxicity. A variety of medications have been involved, including sulfasalazine, cyclosporine, thiopurines, methotrexate as well as certolizumab, adalimumab and infliximab as the biologic agents. Influenza-like symptoms and increased liver enzymes in hepatotoxicity condition usually resolve after the drug treatment has been discontinued²⁷.

2.1.5 Hepatic Steatosis

The most hepatobiliary complication of IBD is hepatic steatosis, or fatty liver²⁸. A group of researchers reported that 35% of the 511 IBD patients had a fatty liver disorder²⁹. In published studies, fatty liver disease has been shown a massive variability in its prevalence (13% to 100%)¹. The level of fatty liver infiltration and the severity of colitis were found to be associated with ulcerative colitis patients³⁰. In IBD patients, the protein deficiency, corticosteroid therapy and chronic

malnutrition, may lead fatty liver condition while the exact causes are somewhat unclear¹.

2.1.6 Hepatic Abscess

The frequency of pyogenic liver abscesses in patients with IBD is slightly greater than in the general population²⁸. Researchers indicated that the loss of barrier integrity of the intestine mucosa may lead to liver damage by an infectious agent through mesenteric veins¹. In addition, portal vein associated- thrombosis can also occur in the portal pyelophlebitis condition³¹.

2.1.7 Portal vein thrombosis

In IBD patients, a thrombosis disorder is commonly observed at the portal vein²⁸. The levels of platelets, fibrinogen and factor V and VIII have been increased in patients with IBD, with antithrombin III being lowered, both of which may raise the risk of thrombosis. The IBD patients who have just undergone abdominal surgery are more likely to develop portal vein thrombosis³². In patients with ulcerative colitis following restorative proctocolectomy, the portal vein thrombosis with a high incidence has been recorded³³.

2.1.8 IBD and Pancreatic manifestations

Concerning IBD-associated pancreatic manifestations in autopsies, in 1950, Ball *et al.*,³⁴ receded on pancreatic features accompanied by UC detected for the first time. In 1956, Chapin *et al.*,³⁵ described that histological changes in the pancreas with regional enteritis were mainly interlobar and periductal fibrosis and swelling of the acinar cells³⁶. Subsequently, in US and Europe, researches on IBD, especially pancreatitis associated CD, have been recorded progressively since the 1970s³⁷⁻³⁹. Pancreatic disorders related with IBD

involve, pancreatic cancer (PC) acute pancreatitis (AP), chronic pancreatitis (CP), increasing of pancreatic enzymes and exocrine pancreatic insufficiency (EPI). Pancreatic manifestations accompanied by IBD has been previously described by Iida *et al.*,⁴⁰ and summarized in Figure 1.

2.1.9 Other diseases associated with IBD

IgG4-associated cholangitis- It was identified in individuals with diseases associated with IgG4 with autoimmune pancreatitis, a relation between ulcerative colitis and cholangitis associated with IgG4 has been observed⁴¹.

Primary biliary cirrhosis- It is a characteristic autoimmune disorder of the hepatic tissue that arises through the progressive degradation of the inflammatory bile ducts and the obliteration of plasma cells and lymphocytes toward the portal way. The literature indicates a correlation between IBD and primary biliary cirrhosis but has not yet been widely recognized⁴².

Cholelithiasis- Gallstones are commonly found in IBD patients. IBD progression and its pathophysiological changes may lead to gallstone development. A substantial correlation has been identified between cholelithiasis abnormality and CD, with 13% and 34% as a prevalence rate⁴³. The link between UC and cholelithiasis is uncertain, because, in ulcerative colitis patients, there does not seem to be a substantially elevated incidence of gallstones relative to the general population²⁹.

Autoimmune Pancreatitis- A correlation between autoimmune pancreatitis and IBD have proposed by some researchers, while to date, the precise association is not well-defined. It has been shown that IBD-associated pancreatitis has similar clinical, morphological, and histological characteristics to the same detected features in autoimmune pancreatitis⁴³.

3. Urinary manifestations of IBD

In the IBD setting, the majority of different urinary complications have been reported in patients with CD, with an incidence rate of 4% -23%. The complications are clearly appearing in people with serious or chronic illness. Urinary conditions can be directly or indirectly associated with the development of the disease⁴⁴.

3.1 Enterourinary Fistulas

These disorders are the most comfortable urinary form in patients with IBD, a colovesical fistulas are the famous type of fistula. While, in CD, gastrointestinal tract-confined fistulas are relatively common, fistulas between the urinary system and the gastrointestinal tract are much less common and severe complications, with an incidence rate of 2%-3.5%⁴⁴. Most patients with a well-known history of IBD are in their 4th or 5th decade. The incidence rate is high in males, due to the position of the adnexa and uterus in women between the intestines and the bladder⁴⁵.

3.2 Obstructive Uropathy

Calculus obstructive uropathy is occurring in 1.9% – 6% of Crohn disease patients, is often ignored urologic complication of Crohn disease⁴⁴. Transmural bowel inflammation may lead to hydroureteronephrosis which followed by ureteral compression, fibrosis, or

encasement and the right collecting ducts usually involved⁴⁶.

3.3 Nephrolithiasis

IBD patients are 10–100 times more susceptible than other patients in the hospital to develop nephrolithiasis, with 12% as an incidence rate in patients with CD⁴⁴. Adults with CD are at higher risk relative to patients with ulcerative colitis than children⁴⁷.

4. IBD and musculoskeletal manifestations

There are multiple musculoskeletal forms of IBD; almost 53% of patients have musculoskeletal system-related pain⁴⁸. The most common sites involved in IBD arthropathy are axial and peripheral⁴⁹.

4.1 IBD-related Spondyloarthropathy

This disease involves psoriatic arthritis, idiopathic ankylosing spondylitis (AS), undifferentiated spondyloarthropathy and reactive arthritis. Spondyloarthropathy associated with IBD is divided into peripheral and axial arthropathy. Symptoms of axial type include back pain, sacroiliitis that triggers inflammation and spondylitis, whereas peripheral arthritis includes self-limiting nondeforming arthritis which develops and decreases with intestinal flares⁵⁰.

4.2 IBD and arthropathy manifestations

Arthropathy is frequent among patients with IBD, and this condition is known as spondylarthritis (SpA). The SpA is classified based on the primary signs, as axial and peripheral⁵¹. A detection of axial SpA is achieved using sacroiliitis radiographic results consistent with signs of little back inflammatory pain. Radiological results of sacroiliitis are significant in around 15%-27% of patients with IBD⁵²⁻⁵⁴, while, advanced AS with syndesmophytes develops in only nearby 3%-10% of patients. Moreover, in CD and AS patients, HLA-B27 is reported in approximately 25%–75% of patients, whereas, is recorded in about 7%–15% in those patients with sacroiliitis, HLA-B27. Positivity of HLA-B27 in IBD patients suggests that these individuals are more vulnerable to develop AS⁵⁵; however, since HLA-B27 positivity in idiopathic AS cases is significantly lower, it cannot be regarded as a diagnostic indicator^{52,56}.

5. IBD and pulmonary manifestations

Black *et al.*,⁵⁷ in their analysis of population research, found that patients with IBD have pulmonary symptoms more often than the general population. Raj *et al.*,⁵⁸ observed a four-fold rise in the incidence of IBD in their patients with airway disease in a 10-year retrospective study. The prevalence of disease-related pulmonary symptoms is highly variable and the symptoms continuum is broad. The pathogenesis of pulmonary findings associated with IBD is uncertain, but may be associated with an embryological origin of the intestinal mucosa and respiratory, a reaction with intestinal epithelium and lung antigen exposure, and/or an intestinal inflammatory mediator⁵⁹.

6. IBD and cardiac manifestations

To date, the link between cardiovascular disorders and IBD has not been completely explained. Researchers have indicated that patients with IBD have a higher risk of myocardial infarction, stroke and cardiovascular mortality in recent years, particularly during the active stage of the disease⁶⁰. The findings of Danish

nationwide cohort study recorded high incidence of cardiac failure in individuals with IBD that was highly associated with active bowel disease periods⁶¹. Other studies have proposed, however, that there is no correlation between increased cardiovascular disease incidence and IBD. Furthermore, a meta-analysis of 11 trials showed no increase in mortality of cardiovascular patients between the control group and IBD group⁶².

7. IBD and ocular system manifestations

According to the findings of the population study in the IBD setting, researchers have reported that the prevalence of ocular extraintestinal symptoms is between 4 % and 12 %⁶³. The most frequent eye investigations include inflammatory disorders affecting various areas of the globe, ranging from episcleritis and conjunctivitis to more serious conditions such as anterior uveitis and scleritis⁶⁴. Vision may be permanently compromised with scleritis, or anterior uveitis, while it is possible to detect unusual orbital extraintestinal manifestations of IBD by imaging tools⁵⁹.

8. IBD Skin diseases manifestations

8.1 Erythema nodosum (EN)

EN is distinguished by the appearance of subcutaneous nodules raised, tender, purple, or violet (1–5 cm in diameter), which makes it simple to diagnose. The extensor outward of the limbs, mostly the frontal areas of tibial, are the most involved areas, and infrequently, the upper limbs or trunk may be involved. Also, EN is related with other symptoms such as weakness and arthralgia. This can be treated by removing the metastatic CD. EN is one the most observed dermatological manifestation in individuals with IBD and is high frequent in women and CD patients (4%-15% CD vs. 3%-10% CD cases)^{65,66}. EN is linked with active IBD in general, however not by its severity⁶⁷. Because of its connection with disease operation, management of the IBD is the cornerstone of therapy. Though, with systemically treatment by corticosteroids may be needed in extreme cases, while treatment with infliximab, azathioprine or adalimumab may be necessary in resistant situations or those with repeated relapses^{68,69}.

8.2 Pyoderma gangrenosum

PG is identified by the formation of a pustule on skin that quickly becomes a violent-edged burrowing ulcer, around 2–20 cm in diameter. PG occurs most often on the legs and close to the stomas, though it can happen anyplace over the body, including genitals. It initially occurs as a solitary or multiple red papule(s), but consequent dermis necrosis contributes to the production of deep gouge chronic ulcerations. For PG, the features of histopathological changes are non-specific, and hence their diagnosis is performed dependent on the characteristic findings of the lesions after ignoring other possible skin disorders depend on the distinctive results of the lesions. Occasionally, to confirm the diagnosis, a biopsy from the lesion may be needed⁵⁵.

CONCLUSION

EIMs are moderately common through the IBD course not only limited to the gut, and in some situations,

these EIM can be much more crippling than bowel disease, which can occur long before IBD is diagnosed. Precise identifying for EIMs and early suitable analysis are imperative to reduce the overall morbidity. Substantial indication recommends that IBD and related extraintestinal illnesses are not isolated disorders but share common pathophysiologic pathways, to date, some of which remain indefinable. In numerous cases, controlling the IBD activity may aid in limiting the EIM; however, the further controlled trials remain the keys to development of the early differentiated diagnosis approach and promising treatment policies.

AUTHOR'S CONTRIBUTION

All authors have worked equally for this work.

ACKNOWLEDGEMENTS

The authors extend their thanks and appreciation to the Qinghai University, China to provide necessary facilities for this work.

CONFLICT OF INTEREST

No conflict of interest associated with this work.

REFERENCES

- Navaneethan U, Shen B. Hepatopancreatobiliary manifestations and complications associated with IBD. *Inflamm Bowel Dis* 2010; 16(9):1598-619. <https://doi.org/10.1002/ibd.21219>
- Torres J, Billioud V, Sachar DB, Peyrin-Biroulet L, Colombel J-F. Ulcerative colitis as a progressive disease: the forgotten evidence. *Inflamm Bowel Dis* 2012; 18(7):1356-63. <https://doi.org/10.1002/ibd.22839>
- Hoivik ML, Moum B, Solberg IC, Henriksen M, Cvancarova M, Bernklev T. Work disability in IBD patients 10 years after disease onset: results from the IBSEN Study. *Gut* 2013; 62(3):368-75. <https://doi.org/10.1136/gutjnl-2012-302311>
- Levesque BG, Sandborn WJ, Ruel J, Feagan BG, Sands BE, Colombel J-F. Converging goals of treatment of IBD from clinical trials and practice. *Gastroenterology* 2015; 148(1):3751.e1. <https://doi.org/10.1053/j.gastro.2014.08.003>
- Dignass A, Lindsay JO, Sturm A, Windsor A, Colombel J-F, Allez M, D'Haens G, D'Hoore A, Mantzaris G, Novacek G. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 2: current management. *J Crohns Colitis* 2012; 6(10):991-1030. <https://doi.org/10.1016/j.crohns.2012.09.002>
- Van Assche G, Dignass A, Bokemeyer B, Danese S, Gionchetti P, Moser G, Beaugerie L, Gomollón F, Häuser W, Herrlinger K. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 3: special situations. *Rev Gastroenterol Mex* 2015; 80(1):74-106. <https://doi.org/10.1016/j.rgm.2014.10.008>
- Truelove S. Corisone in ulcerative colitis. Final report on a therapeutic trial. *Br Med J* 1955 29; 2(4947):1041-8. <https://doi.org/10.1136/bmj.2.4947.1041>
- Colombel JF, Rutgeerts P, Reinisch W, Esser D, Wang Y, Lang Y, Marano CW, Strauss R, Oddens BJ, Feagan BG. Early mucosal healing with infliximab is associated with improved long-term clinical outcomes in ulcerative colitis. *Gastroenterol* 2011; 141(4):1194-201. <https://doi.org/10.1053/j.gastro.2011.06.054>
- Lewis JD, Chuai S, Nessel L, Lichtenstein GR, Aberra FN, Ellenberg JH. Use of the noninvasive components of

- the Mayo score to assess clinical response in ulcerative colitis. *Inflamm Bowel Dis* 2008; 14(12):1660-6.
<https://doi.org/10.1002/ibd.20520>
10. Peyrin-Biroulet L, Sandborn W, Sands B, Reinisch W, Bemelman W, Bryant R, d'Haens G, Dotan I, Dubinsky M, Feagan B. Selecting therapeutic targets in IBD (STRIDE): determining therapeutic goals for treat-to-target. *Am J Gastroenterol* 2015; 110(9):1324-38.
<https://doi.org/10.1038/ajg.2015.233>
 11. Yoon JY, Park SJ, Hong SP, Kim TI, Kim WH, Cheon JH. Correlations of C-reactive protein levels and erythrocyte sedimentation rates with endoscopic activity indices in patients with ulcerative colitis. *Dig Dis Sci* 2014; 59(4):829-37.
<https://doi.org/10.1007/s10620-013-2907-3>
 12. Dignass A, Eliakim R, Magro F, Maaser C, Chowers Y, Geboes K, Mantzaris G, Reinisch W, Colombel J-F, Vermeire S. Second European evidence-based consensus on the diagnosis and management of ulcerative colitis part 1: definitions and diagnosis. *J Crohns Colitis* 2012; 6(10):965-90.
<https://doi.org/10.1016/j.crohns.2012.09.003>
 13. Bessissow T, Lemmens B, Ferrante M, Bisschops R, Van Steen K, Geboes K, Van Assche G, Vermeire S, Rutgeerts P, De Hertogh G. Prognostic value of serologic and histologic markers on clinical relapse in ulcerative colitis patients with mucosal healing. *Am J Gastroenterol* 2012; 107(11):1684-92.
<https://doi.org/10.1038/ajg.2012.301>
 14. Peyrin-Biroulet L, Bressenot A, Kampman W. Histologic remission: the ultimate therapeutic goal in ulcerative colitis? *Clin Gastroenterol Hepatol* 2014; 12(6):929-34.e2.
<https://doi.org/10.1016/j.cgh.2013.07.022>
 15. Bitton A, Peppercorn MA, Antonioli DA, Niles JL, Shah S, Bousvaros A, Ransil B, Wild G, Cohen A, Edwardes MDD. Clinical, biological, and histologic parameters as predictors of relapse in ulcerative colitis. *Saudi J Gastroenterol* 2011; 17(3):194-8.
<https://doi.org/10.4103/1319-3767.80383>
 16. Bryant RV, Burger DC, Delo J, Walsh AJ, Thomas S, von Herbay A, Buchel OC, White L, Brain O, Keshav S. Beyond endoscopic mucosal healing in UC: histological remission better predicts corticosteroid use and hospitalisation over 6 years of follow-up. *Gut* 2016; 65(3):408-14.
<https://doi.org/10.1136/gutjnl-2015-309598>
 17. Kornbluth A, Sachar DB, Gastroenterology PP Cot A Co. Ulcerative colitis practice guidelines in adults: American college of gastroenterology, practice parameters committee. *Am J Gastroenterol* 2010; 105(3):501-23; quiz 524. <https://doi.org/10.1038/ajg.2009.727>
 18. Sawczenko A, Sandhu B. Presenting features of IBD in Great Britain and Ireland. *Arch Dis Child* 2003; 88(11):995-1000. <https://doi.org/10.1136/adc.88.11.995>
 19. Cummings JF, Keshav S, Travis SP. Medical management of CD. *BMJ* 2008 10; 336(7652):1062-6.
<https://doi.org/10.1136/bmj.39547.603218.AE>
 20. Bernstein CN, Blanchard JF, Rawsthorne P, Yu N. The prevalence of extraintestinal diseases in IBD: a population-based study. *Am J Gastroenterol* 2001; 96(4):1116-22.
<https://doi.org/10.1111/j.1572-0241.2001.03756.x>
 21. Olpin JD, Sjöberg BP, Stilwill SE, Jensen LE, Rezvani M, Shaaban AM. Beyond the bowel: extraintestinal manifestations of IBD. *Radiographics* 2017; 37:1135-60.
<https://doi.org/10.1148/rf.2017160121>
 22. Zhang W, Nuki G, Moskowitz R, Abramson S, Altman RD, Arden N, Bierma-Zeinstra S, Brandt K, Croft P, Doherty M. OARSIS recommendations for the management of hip and knee osteoarthritis: part III: Changes in evidence following systematic cumulative update of research published through January 2009. *Osteoarthritis Cartilage* 2010; 18(4):476-99.
<https://doi.org/10.1016/j.joca.2010.01.013>
 23. Smith MP, Loe RH: Sclerosing cholangitis. review of recent case reports and associated diseases and four new cases. *Am J Surg* 1996; 110:239-46.
[https://doi.org/10.1016/0002-9610\(65\)90018-8](https://doi.org/10.1016/0002-9610(65)90018-8)
 24. Floreani A, Rizzotto ER, Ferrara F, Carderi I, Caroli D, Blasone L, Baldo V. Clinical course and outcome of autoimmune hepatitis/primary sclerosing cholangitis overlap syndrome. *Am J Gastroenterol* 2005; 100(7):1516-22.
<https://doi.org/10.1111/j.1572-0241.2005.41841.x>
 25. Broome U, Olsson R, Lööf L, Bodemar G, Hultcrantz R, Danielsson A, Prytz H, Sandberg-Gertzen H, Wallerstedt S, Lindberg G. Natural history and prognostic factors in 305 Swedish patients with primary sclerosing cholangitis. *Gut* 1996; 38(4):610-5.
<https://doi.org/10.1136/gut.38.4.610>
 26. Walker SL, McCormick PA. Diagnosing cholangiocarcinoma in primary sclerosing cholangitis: an "evidence based radiology" review. *Abdom Imaging* 2008; 33:14-7. <https://doi.org/10.1007/s00261-007-9314-1>
 27. Roenigk HH, Auerbach R, Maibach H, Weinstein G, Lebwohl M. Methotrexate in psoriasis: consensus conference. *J Am Acad Dermatol* 1998; 38(3):478-85.
[https://doi.org/10.1016/s0190-9622\(98\)70508-0](https://doi.org/10.1016/s0190-9622(98)70508-0)
 28. Restellini S, Chazouillères O, Frossard JL. Hepatic manifestations of IBDs. *Liver Int* 2017; 37(4):475-489.
<https://doi.org/10.1111/liv.13265>
 29. Bargiggia S, Maconi G, Elli M, Molteni P, Ardizzone S, Parente F, Todaro I, Greco S, Manzionna G, Porro GB. Sonographic prevalence of liver steatosis and biliary tract stones in patients with IBD: study of 511 subjects at a single center. *J Clin Gastroenterol* 2003; 36(5):417-20.
<https://doi.org/10.1097/00004836-200305000-00012>
 30. Riegler G, D'INCÀ R, Sturmiolo G, Corrao G, Blanco CDV, Di Leo V, Carratù R, Ingrosso M, Pelli M, Morini S. Hepatobiliary alterations in patients with IBD: a multicenter study. *Scand J Gastroenterol* 1998; 33(1):93-8. <https://doi.org/10.1080/00365529850166275>
 31. Vakil N, Hayne G, Sharma A, Hardy DJ, Slutsky A. Liver abscess in CD. *American J Gastroenterol* 1994; 89. PMID: 8017370
 32. Jackson L, O'Gorman P, O'Connell J, Cronin C, Cotter K, Shanahan F. Thrombosis in IBD: clinical setting, procoagulant profile and factor V Leiden. *QJM: An Int J Med* 1997; 90(3):183-8.
<https://doi.org/10.1093/qjmed/90.3.183>
 33. Baker ME, Remzi F, Einstein D, Oncel M, Herts B, Remer E, Fazio V. CT depiction of portal vein thrombi after creation of ileal pouch-anal anastomosis. *Radiol* 2003; 227(1):73-9.
<https://doi.org/10.1148/radiol.2271020032>
 34. Ball WP. Pancreatic lesions associated with chronic ulcerative colitis. *Arch Pathol (Chic)*. 1950; 50(3):347-58. PMID: 15433704
 35. Chapin LE, Scudamore HH, Baggenstoss AH, Bagen JA. Regional enteritis: associated visceral changes. *Gastroenterol* 1956; 30(3):404-15.
[https://doi.org/10.1016/S0016-5085\(56\)80104-2](https://doi.org/10.1016/S0016-5085(56)80104-2)
 36. Frey CF. Acute pancreatitis as a complication of ulcerative colitis and collagen disease. *Univ Mich Med Cent J: University of Michigan Medical Center J* 1967; 33(1):18-21. PMID: 6037879
 37. Legge DA, Hoffman HN, Carlson HC. Pancreatitis as a complication of regional enteritis of the duodenum. *Gastroenterol* 1971; 61(6):834-7. PMID: 5125685
 38. Meltzer SJ, Korelitz BI. Pancreatitis and duodenopancreatic reflux in CD. Case report and review of the literature. *J Clin Gastroenterol* 1988; 10(5):555-8.
<https://doi.org/10.1097/00004836-198810000-00016>
 39. Seyrig J-A, Jian R, Modigliani R, Golfain D, Florent C, Messing B, Bitoun A. Idiopathic pancreatitis associated with IBD. *Dig Dis Sci* 1985; 30(12):1121-6.
<https://doi.org/10.1007/BF01314044>

40. Iida T, Wagatsuma K, Hirayama D, Yokoyama Y, Nakase H. The Etiology of Pancreatic Manifestations in Patients with IBD. *J Clin Med* 2019 26; 8(7):916. <https://doi.org/10.3390/jcm8070916>
41. Dastis SN, Latinne D, Sempoux C, Geubel AP. Ulcerative colitis associated with IgG4 cholangitis: similar features in two HLA identical siblings. *J Hepatol* 2009; 51(3):601-5. <https://doi.org/10.1016/j.jhep.2009.05.032>
42. Tada F, Abe M, Nunoi H, Azemoto N, Mashiba T, Furukawa S, Kumagi T, Murakami H, Ikeda Y, Matsuura B. Ulcerative colitis complicated with primary biliary cirrhosis. *I Intern Med* 2011; 50(20):2323-7. <https://doi.org/10.2169/internalmedicine.50.5919>
43. Navaneethan U, Shen B. Hepatopancreatobiliary manifestations and complications associated with IBD. *Inflamm Bowel Dis* 2010; 16(9):1598-619. <https://doi.org/10.1002/ibd.21219>
44. Tonolini M, Villa C, Campari A, Ravelli A, Bianco R, Cornalba G. Common and unusual urogenital CD complications: spectrum of cross-sectional imaging findings. *Abdom Imaging* 2013; 38(1):32-41. <https://doi.org/10.1007/s00261-012-9876-4>
45. Talamini M, Broe P, Cameron JL. Urinary fistulas in CD. *Surg Gynecol Obstet* 1982; 154(4):553-6. PMID: 7064089
46. Ruffolo C, Angriman I, Scarpa M, Polese L, Pagano D, Barollo M, Bertin M, D'Amico DF. Urologic complications in CD: suspicion criteria. *Hepatogastroenterol* 2006; 53(69):357-60. PMID: 16795972
47. McLeod RS, Churchill DN. Urolithiasis complicating IBD. *J Urol* 1992; 148(3 Pt 2):974-8. [https://doi.org/10.1016/s0022-5347\(17\)36794-0](https://doi.org/10.1016/s0022-5347(17)36794-0)
48. Levine JS, Burakoff R. Extraintestinal manifestations of IBD. *Gastroenterol Hepatol (NY)* 2011; 7(4):235-41.
49. Arvikar SL, Fisher MC. IBD associated arthropathy. *Curr Rev Musculoskelet Med* 2011; 4(3):123-31. <https://doi.org/10.1007/s12178-011-9085-8>
50. Brakenhoff LK, van der Heijde DM, Hommes DW, Huizinga TW, Fidler HH. The joint—gut axis in IBDs. *J Crohns Colitis* 2010; 4(3):257-68. <https://doi.org/10.1016/j.crohns.2009.11.005>
51. Rudwaleit M, Van Der Heijde D, Landewé R, Listing J, Akkoc N, Brandt J, Braun J, Chou C, Collantes-Estevez E, Dougados M. The development of Assessment of SpondyloArthritis international Society classification criteria for axial spondyloarthritis (part II): validation and final selection. *Ann Rheum Dis* 2009; 68(6):777-83. <https://doi.org/10.1136/ard.2009.108233>
52. Chan J, Sari I, Salonen D, Silverberg MS, Haroon N, Inman RD. Prevalence of sacroiliitis in IBD using a standardized computed tomography scoring system. *Arthritis Care Res (Hoboken)* 2018; 70(5):807-810. <https://doi.org/10.1002/acr.23323>
53. Leclerc JS, Lux G, Rat A, Laurent V, Blum A, Chary VI, Peyrin BL, Loeuille D. The prevalence of inflammatory sacroiliitis assessed on magnetic resonance imaging of IBD: a retrospective study performed on 186 patients. *Aliment Pharmacol Ther* 2014; 39(9):957-62. <https://doi.org/10.1111/apt.12680>
54. Peeters H, Vander Cruyssen B, Mielants H, De Vlam K, Vermeire S, Louis E, Rutgeerts P, Belaiche J, De Vos M. Clinical and genetic factors associated with sacroiliitis in CD. *J Gastroenterol Hepatol* 2008; 23(1):132-7. <https://doi.org/10.1111/j.1440-1746.2007.05108.x>
55. Harbord M, Annesse V, Vavricka SR, Allez M, Barreiro-de Acosta M, Boberg KM, Burisch J, De Vos M, De Vries A-M, Dick AD. The first European evidence-based consensus on extra-intestinal manifestations in IBD. *J Crohns Colitis* 2016; 10(3):239-54. <https://doi.org/10.1093/ecco-jcc/jjv213>
56. Orchard T, Holt H, Bradbury L, Hammersma J, McNally E, Jewell D, Wordsworth B. The prevalence, clinical features and association of HLA-B27 in sacroiliitis associated with established CD. *Aliment Pharmacol Ther* 2009; 29(2):193-7. <https://doi.org/10.1111/j.1365-2036.2008.03868.x>
57. Black H, Mendoza M, Murin S. Thoracic manifestations of IBD. *Chest* 2007; 131(2):524-32. <https://doi.org/10.1378/chest.06-1074>
58. Raj A, Birring S, Green R, Grant A, de Caestecker J, Pavord I. Prevalence of IBD in patients with airways disease. *Respir Med* 2008; 102(5):780-5. <https://doi.org/10.1016/j.rmed.2007.08.014>
59. Katsanos A, Asproudis I, Katsanos KH, Dastiridou AI, Aspiotis M, Tsianos EV. Orbital and optic nerve complications of IBD. *J Crohns Colitis* 2013; 7(9):683-93. <https://doi.org/10.1016/j.crohns.2012.09.020>
60. Kristensen SL, Ahlehoff O, Lindhardsen J, Erichsen R, Jensen GV, Torp-Pedersen C, Nielsen OH, Gislason GH, Hansen PR. Disease activity in IBD is associated with increased risk of myocardial infarction, stroke and cardiovascular death—a Danish nationwide cohort study. *PLoS One*. 2013; 8(2):e56944. <https://doi.org/10.1371/journal.pone.0056944>
61. Kristensen SL, Ahlehoff O, Lindhardsen J, Erichsen R, Lamberts M, Khalid U, Nielsen OH, Torp-Pedersen C, Gislason GH, Hansen PR. IBD is associated with an increased risk of hospitalization for heart failure: a Danish Nationwide Cohort study. *Circ Heart Fail* 2014; 7(5):717-22. <https://doi.org/10.1161/CIRCHEARTFAILURE.114.001152>
62. Dorn SD, Sandler RS. IBD is not a risk factor for cardiovascular disease mortality: results from a systematic review and meta-analysis. *Am J Gastroenterol* 2007; 102(3):662-7. <https://doi.org/10.1111/j.1572-0241.2006.01018.x>
63. Das KM. Relationship of extraintestinal involvements in IBD (new insights into autoimmune pathogenesis). *Dig Dis Sci* 1999; 44(1):1-13. <https://doi.org/10.1023/a:1026629528233>
64. Rothfuss KS, Stange EF, Herrlinger KR. Extraintestinal manifestations and complications in IBDs. *World journal of gastroenterology: World J Gastroenterol* 2006 14; 12(30):4819-31. <https://doi.org/10.3748/wjg.v12.i30.481>
65. Chowaniec M, Starba A, Wiland P. Erythema nodosum—review of the literature. *Reumatologia* 2016; 54(2):79-82. <https://doi.org/10.5114/reum.2016.60217>
66. Freeman HJ. Erythema nodosum and pyoderma gangrenosum in 50 patients with CD. *Can J Gastroenterol* 2005; 19(10):603-6. <https://doi.org/10.1155/2005/323914>
67. Trost L, McDonnell J. Important cutaneous manifestations of IBD. *Postgrad Med J* 2005; 81(959):5805. <https://doi.org/10.1136/pgmj.2004.031633>
68. Clayton T, Walker B, Stables G. Treatment of chronic erythema nodosum with infliximab. *Clin Exp Dermatol* 2006; 31(6):823-4. <https://doi.org/10.1111/j.1365-2230.2006.02221.x>
69. Ortego CN, Callejas RJ, Sanchez CD, Caballero MT. Refractory chronic erythema nodosum successfully treated with adalimumab. *J Eur Acad Dermatol Venereol* 2007; 21(3):408-10. <https://doi.org/10.1111/j.1468-3083.2006.01893.x>